



CRISPR/Cas9 genome editing tool for rice crop improvement

Selvakumar Gurunathan^{1*}, Bharathi Raja Ramadoss², Ambika Singaram¹, Dhanarajan Arulbalachandran³

Received: 7 February 2023 / Accepted: 14 April 2023 / Published: 30 June 2023

¹Department of Crop Improvement, SRM College of Agricultural Sciences, SRM Institute of Science and Technology, Vendhar Nagar, Baburayanpettai, Tamil Nadu, India.

²Bioriginal Food and Science Corporation, Saskatoon, SK S7J 0R1, Canada.

³Division of Crop Molecular Breeding and Stress Physiology, Department of Botany, School of Life Sciences, Periyar University, Periyar Palkalai Nagar, Salem, Tamil Nadu, India.

*Correspondence

Selvakumar Gurunathan

selvakumaragri@gmail.com; selvakug@srmist.edu.in

Food crop yield, quality, and tolerance mechanisms to biotic and abiotic factors are important aspects that contribute to food security. To feed about 10 billion people by 2050, high yielding climate-resilient rice cultivars with good grain quality must be created more quickly. Yield and quality, along with stress tolerance traits of the rice crop, have been improved by adopting various methods. Among these, in recent years, the yield of the crop has been improved marginally by utilizing conventional breeding methods. Mutation breeding is an important pathway that has created many novel variations and contributed towards isolating new high yielding genotypes in the rice crop. Forward and reverse genetic protocols have been engaged for the identification of genomic variants in conventional mutation breeding to characterize the novel variants to convert as functional markers for the development of new improved varieties. Generation of desired mutations in the desirable region of the genome of the crops is highly tedious through conventional breeding methods such as random mutagenesis since the gene manipulations happen randomly while the mutagenesis is done using physical and chemical mutagens. Also, it requires large mutant plant populations to isolate the desired mutants and mutations. The advancement of CRISPR/Cas9 genome editing technology rapidly replaces conventional random mutagenesis technologies, has the ability to multiplex genome editing to create novel variations for crop improvement programs, and reduces the time duration required for trait-based crop improvement programs. In this review, significant gene manipulations employed through CRISPR/Cas9 for rice crop improvement in terms of yield and biotic and abiotic stress tolerance are discussed.

Key words: CRISPR/Cas9, crop improvement, mutagenesis, mutations, genome editing, rice

Introduction

Currently, the rice crop feeds and ensures food security for nearly half of the world's population. According to the FAO (Romero & Gatica-Arias, 2019), the world's projected population of ten billion will necessitate a 74% increase in food demand by 2050 due to the alarming population incremental phenomenon. More specifically, a 40% increase in rice consumption is anticipated (Milovanovic & Smutka, 2017). Even though there are many high-yielding varieties and hybrids of rice available right now, there is still a need to develop high-yielding rice varieties that can adapt to changing environments (Clarke & Zhang, 2013) in order to feed the world's growing population. Over the past few years, numerous conventional methods have been employed to increase rice yield (Khush, 2003; Brennan & Malabayabas, 2011; Peng et al., 2008; Denardin et al., 2019) in the face of the strain (Blum, 2009; Seo et al., 2020; Tripathi et al., 2012; Mackill et al., 2010). Genetic engineering methodologies are also used to improve the grain quality of rice crops (Ryoo et al., 2007; Zhang et al., 2013) along with salt stress (Yue et al., 2020) and drought (Li et al., 2021) tolerance. Conventional crop improvement methods such as in random mutagenesis, different types of physical and chemical mutagens are utilized to create new variation in the genome of crop plants. The identification of desired genotypes contributes to crop improvement after the creation of the mutant population. The maintenance

screening, and development of a random mutagenized population are extremely time-consuming and laborious. While attempting to improve related traits in crops through specific genome mutations, sometimes fail to achieve the desired genotypes.

The development of technology for manipulating genes, the combination of CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats) and the CRISPR-associated protein Cas9 is an excellent tool which is capable for generating novel genomic variations in targeted regions (Zafar et al., 2020) and also it is comparatively superior than TALENs (Zafar et al., 2020). CRISPR/Cas9 technology rapidly replaces random mutagenesis crop improvement programs and is extremely useful for multiplex genome editing to create novel variations. CRISPR/Cas9 technology facilitates trait-based crop development by causing double strand breaks in the DNA sequence in the targeted location. In terms of crop improvement, CRISPR/Cas9 is the unavoidable tool. In particular, the application of this technology to rice (*Oryza sativa*) will result in the creation of novel genotypes that will improve yield and biotic and abiotic stress tolerance. Since rice serves as a model plant for functional genomic analysis, the utilization of CRISPR/Cas9 paves the way for the enhancement of related traits in other cereal crops. Enhanced specificity of CRISPR/Cas9 on the genome editing in the DNA sequence is being utilized for the identification of the function of the genes and molecular breeding strategies. This technology requires very short time to create the new variations to develop new cultivars and improve the existing cultivars with improved performance in terms of yield and other related traits for the efficient crop production.

Yield improvement

Grain yield is a complex trait yet an essential and inevitable strives of rice crop improvement programmes. Yielding ability of the rice is controlled by polygenes and majorly contributed by grain numbers/panicle, panicle number/plant and 1000 grain weight (Xing & Zhang, 2010). These days, rice assortments utilizing genome-altering innovations contain new genotypes of yield-related qualities. Shan et al., (2014) reported the potential of CRISPR/Cas9 framework to increase the grain yield of the rice crop. CRISPR/Cas9 is used to modify the genes viz., *Gn1a*, *DEP1*, *GS3*, and *IPA1* which are regulatory genes for the traits, the architecture of the plant and panicle, number of grains per panicle and size of the grains. Various research undertakings have been made to additionally foster grain yield using CRISPR-Cas9 (Sedeek et al., 2019). Grain size is one of the most important parameters influencing rice grain quality and yield potential. It has sparked considerable interest among molecular biologists and breeders for yield improvement (Wang et al., 2015). Wang et al., (2022) targeted miR396 site present in the CS2 gene which regulates a few essential traits such as seed shattering, grain size, grain quality and nitrogen use efficiency and abiotic stress response and resulted in a gain of function in the GS2^E mutant. The mutant showed multiple beneficial trait expressions on grain size and yield; also, the thousand grain weight was increased by 23.5%, and consequently, the yield was increased by 10.4%.

Biotic and abiotic stress

Targeting regulators and genes which are responsible for the biotic and abiotic resistance or tolerances in the crop plants for the modification or to generate the allelic forms could be a best way for the development of the biotic and abiotic resistance or tolerances plants. Since rice genomes are available with excellent sequence quality increases the possibility of selection and modification of the specific genes.

Table 1. Recent applications of CRISPR/Cas9 technology on biotic stress tolerance enhancement in rice

Diseases	Targeted gene	Citation
Blast	<i>OsDjA2</i> and <i>OsERF104</i>	Távora et al., 2022
Blast	<i>Bsr-d1</i> , <i>Pi21</i> and <i>ERF922</i>	Zhou et al., 2022
Bacterial blight	<i>Os8N3</i>	Kim et al., 2019
Bacterial blight	<i>EBEs</i> (Promotor regions of <i>OsSWEET14</i>)	Zafar et al., 2020
Bacterial blight	<i>OsSWEET14</i>	Zeng et al., 2020

Currently, CRISPR/Cas9 genome editing technology hastens crop development programmes regarding biotic and abiotic stress resistance or tolerances genotypes (Table 1). The main challenges for crop plants during both vegetative

and reproductive stages are biotic and abiotic stress to express maximum productivity. Blast is the most adverse disease in rice crop caused by *Pyricularia oryzae* (syn. *Magnaporthe oryzae*) one and reducing crop yield worldwide (Jain et al. 2017). Cultivation of rice varieties with multiple key resistant genes for blast disease is the most commonly used method to manage this disease.

Cultivation of rice resistant cultivars with single or multiple key resistance (R) genes is the most commonly used and most environmentally friendly approach to control *P. oryzae* infection. Távora et al., 2022 knocked out *OsDjA2* and *OsERF104* genes and explained about the level of disease resistance in the mutant Nipponbare rice plants which showed reduced number of blast lesions and reduced percentage of diseased leaf area compared to control plants. A *indica* TGMS line (LK638S) used for the generation of mutations in the genes *Bsr-d1*, *Pi21* and *ERF922* and created single as well as triple mutants through CRISPR/Cas9 (Zhou et al., 2022). The results of these experiments revealed that *ERF922* mutants possessed strongest resistance nature against blast disease. The gene *Os8N3* is otherwise known as *OsSWEET14*, which is functioning as sugar transporter. Kim et al. (2019) targeted EBEs of *Os8N3* for knocked out this gene and conferred the bacterial leaf blight disease resistance in the Kitaake (*Oryza sativa* L. ssp. *Japonica*) cultivar. In Basmati varieties, bacterial blight caused by *Xanthomonas oryzae* pv. *Oryzae* led to considerable yield losses. Zafar et al., 2020 targeted the four EBSs which are residing in the promotor region of *OsSWEET14* genes for the CRISPR Cas9 editing and created TALEs (*AvrXa7*, *PthXo3*, and *TalF*). The deletions in the EBE of *AvrXa7* revealed the resistance against locally prevailing *Xoo* strains. Zeng et al. (2020) utilized CRISPR/Cas9 and created different mutant allelic natures of *OsSWEET14* genes in the rice cv. Zhonghua 11 (*CR-S14*) and conferred broad spectrum disease resistance against Asian and African *Xoo* strains. At the same time, the yielding potential of the genotypes are not disturbed due to the disruption in the *OsSWEET14*, but the plant height was increased.

Abiotic stress-tolerant rice cultivars must be developed if rice production is to be sustained in the face of rising salt-affected areas, diminishing freshwater supplies, and climate change. EMS mutants and transgenics have been used to validate the function of several rice genes. Frequently, a large number of these positive alleles are not accessible to rice which must be essentially developed. It will be used for introgression breeding to resolve the abiotic stress susceptibility of improved rice cultivars. Salinity is one of the most significant abiotic stresses affecting rice production worldwide. Cultivating salinity-tolerant cultivars is the most environmentally friendly and cost-effective method for managing salinity. For target-site genome editing, CRISPR/Cas9 systems have become increasingly popular in recent years to develop the salinity tolerance crop varieties. CRISPR/Cas9 employed targeted mutagenesis in the *OsRR22* was improved the salinity tolerance in the japonica rice cultivar WPB106 (Zhang et al., 2019). Kim et al. (2023) paved the way of generating targeted mutagenesis in the gene *OsPUB7* through CRISPR/Cas9 to develop drought tolerance and abiotic stress rice in the future. Alam et al. (2022) reported that knocking out the *OsBHLH024* transcription factor conferred salinity tolerance. A deletion mutation in mutant A91 caused an increased level of expression in the genes *OsHKT1*, *OsHKT3*, *OsHAK7*, and *OsSOS1* under salt stress.

Using CRISPR-Cas9 gene editing in *indica* rice cv. MTU1010, generated mutant alleles of the drought and salt tolerance (*DST*) gene (Santosh Kumar et al., 2020). Two distinct gRNAs have been utilized to generate *DST* mutant alleles. The function of the *DST* protein might be affected the protein–protein interaction functions. Loss of function mutation in the *DST* mutants revealed decreased stomatal density which is caused by the downregulation of the stomatal development genes *SPCH1*, *MUTE*, and *ICE1*. The mutants exhibited moderate osmotic stress tolerance and elevated salt stress tolerance. Through improving the morphology and rolling trait in the leaf of the rice will help to get sustainable crop yield under the water deficit conditions. Mutation created in the *Semi-rolled leaf1,2* (*SRL1* and *SRL2*) genes generated the leaf rolling traits during the water deficit stress in the plants (Liao et al., 2019). The results revealed that the chlorophyll content, transpiration rate, stomatal conductance, vascular bundles (VB), stomatal number, and agronomic traits of homozygous mutants were all reduced, while the number of panicles and bulliform cells (BCs) were increased. The hybridization of mutant and its restorer showed a phenotype of semi-rolled leaves, more panicles, more grains per panicle, and more yield per plant.

Zeng et al. (2020) utilized the CRISPR/Cas9 targeted mutagenesis in the genes *OsPIN5b* (panicle length), *GS3* (grain size) and *OsMYB30* (cold tolerance). This experiment reported the improved yielding ability of the rice crop along with improved tolerance to cold stress. Park et al., 2022 adopted CRISPR/Cas9 to edit the *Oryza sativa* *Senescence-associated protein* (*OsSAP*) and it is otherwise called as drought induced genes and concluded that it is an efficient

and novel tool to generate novel variation in the genes in the targeted region in rice crop. CRISPR/Cas9 mediated *afp1* mutants were analysed (Tianshun et al., 2021) and the results showed that Plant height and seed setting rate were lower in *afp1* mutants under normal conditions, but number of tillers/plant and length of the panicle were significantly increased. Interestingly, single plant yield of the mutant plant was showed significant variations. The *afp1* mutants had a lower ABA sensitivity and decreased water loss rate while compared with parent plant. The resistance to heat, drought, and osmotic stress were significantly increased.

Conclusion

Though many conventional crop breeding methodologies contribute to food security, increasing crop productivity is an important issue to address in the given scenario of increasing population and climatic fluctuations. A rapid method is needed to study the gene functions and generation of new variations in crop plants to develop new cultivars with improved yields as well as improved biotic and abiotic stress tolerances. The advent of genome editing tools, especially CRISPR/Cas, hastens the speed of the aforementioned objectives. It helps to edit the targeted genomic regions for the trait specific crop improvement program in a short time. From a scientific perspective, the label-free transgenic mutants and varieties produced by CRISPR/Cas are identical to those produced by natural mutation or conventional mutagenesis. Hence, the CRISPR/Cas technology could be exploited further for the development of new and improved varieties with the adaptation of unpredictable climatic changes to face the food security crisis in the future.

Author contributions

SG: Conceptualized and developed this manuscript. BRR, AS & DA: Edited this manuscript and fine-tuned it.

Competing interests

The authors have declared that no conflict of interest exists.

Ethics approval

Not applicable

References

- Alam, M. S., Kong, J., Tao, R., Ahmed, T., Alamin, M., Alotaibi, S. S., ... & Xu, J. H. (2022). CRISPR/Cas9 mediated knockout of the OsbHLH024 transcription factor improves salt stress resistance in rice (*Oryza sativa* L.). *Plants*, 11(9), 1184.
- Blum, A. (2009). Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. *Field crops research*, 112(2-3), 119-123.
- Brennan, J. P., & Malabayabas, A. (2011). International Rice Research Institute's contribution to rice varietal yield improvement in South-East Asia.
- Clarke, J. L., & Zhang, P. (2013). Plant biotechnology for food security and bioeconomy. *Plant Molecular Biology*, 83, 1-3.
- Denardin, L. G. D. O., Carmona, F. D. C., Veloso, M. G., Martins, A. P., de Freitas, T. F. S., Carlos, F. S., ... & Anghinoni, I. (2019). No-tillage increases irrigated rice yield through soil quality improvement along time. *Soil and Tillage Research*, 186, 64-69.

- Jain, P., Singh, P. K., Kapoor, R., Khanna, A., Solanke, A. U., Krishnan, S. G., ... & Sharma, T. R. (2017). Understanding host-pathogen interactions with expression profiling of NILs carrying rice-blast resistance Pi9 gene. *Frontiers in Plant Science*, 8, 93.
- Khush, G. (2003). Productivity improvements in rice. *Nutrition reviews*, 61(suppl_6), S114-S116.
- Kim, M. S., Ko, S. R., Jung, Y. J., Kang, K. K., Lee, Y. J., & Cho, Y. G. (2023). Knockout Mutants of OsPUB7 Generated Using CRISPR/Cas9 Revealed Abiotic Stress Tolerance in Rice. *International Journal of Molecular Sciences*, 24(6), 5338.
- Kim, Y. A., Moon, H., & Park, C. J. (2019). CRISPR/Cas9-targeted mutagenesis of Os8N3 in rice to confer resistance to *Xanthomonas oryzae* pv. *oryzae*. *Rice*, 12(1), 1-13.
- Li, J., Zhang, M., Yang, L., Mao, X., Li, J., Li, L., ... & Zou, D. (2021). OsADR3 increases drought stress tolerance by inducing antioxidant defense mechanisms and regulating OsGPX1 in rice (*Oryza sativa* L.). *The Crop Journal*, 9(5), 1003-1017.
- Liao, S., Qin, X., Luo, L., Han, Y., Wang, X., Usman, B., ... & Li, R. (2019). CRISPR/Cas9-induced mutagenesis of semi-rolled leaf1, 2 confers curled leaf phenotype and drought tolerance by influencing protein expression patterns and ROS scavenging in rice (*Oryza sativa* L.). *Agronomy*, 9(11), 728.
- Mackill, D. J., Ismail, A. M., Pamplona, A. M., Sanchez, D. L., Carandang, J. J., & Septiningsih, E. M. (2010). Stress tolerant rice varieties for adaptation to a changing climate. *Crop, Environment & Bioinformatics*, 7, 250-259.
- Milovanovic, V., & Smutka, L. (2017). Asian countries in the global rice market. *ACTA Universitatis agriculturae et silviculturae mendelianae Brunensis*, 65(2), 679-688.
- Park, J. R., Kim, E. G., Jang, Y. H., Jan, R., Farooq, M., Ubaidillah, M., & Kim, K. M. (2022). Applications of CRISPR/Cas9 as new strategies for short breeding to drought gene in rice. *Frontiers in Plant Science*, 13.
- Peng, S., Khush, G. S., Virk, P., Tang, Q., & Zou, Y. (2008). Progress in ideotype breeding to increase rice yield potential. *Field Crops Research*, 108(1), 32-38.
- Romero, F. M., & Gatica-Arias, A. (2019). CRISPR/Cas9: development and application in rice breeding. *Rice Science*, 26(5), 265-281.
- Ryoo, N., Yu, C., Park, C. S., Baik, M. Y., Park, I. M., Cho, M. H., ... & Jeon, J. S. (2007). Knockout of a starch synthase gene OsSSIIIa/Flo5 causes white-core floury endosperm in rice (*Oryza sativa* L.). *Plant cell reports*, 26, 1083-1095.
- Santosh Kumar, V. V., Verma, R. K., Yadav, S. K., Yadav, P., Watts, A., Rao, M. V., & Chinnusamy, V. (2020). CRISPR-Cas9 mediated genome editing of drought and salt tolerance (OsDST) gene in indica mega rice cultivar MTU1010. *Physiology and Molecular Biology of Plants*, 26, 1099-1110.
- Seo, D. H., Seomun, S., Choi, Y. D., & Jang, G. (2020). Root development and stress tolerance in rice: the key to improving stress tolerance without yield penalties. *International Journal of Molecular Sciences*, 21(5), 1807.
- Shan, Q., Wang, Y., Li, J., & Gao, C. (2014). Genome editing in rice and wheat using the CRISPR/Cas system. *Nature protocols*, 9(10), 2395-2410.
- Távora, F. T., Meunier, A. C., Vernet, A., Portefaix, M., Milazzo, J., Adreit, H., ... & Mehta, A. (2022). CRISPR/Cas9-Targeted knockout of rice susceptibility genes OsDjA2 and OsERF104 reveals alternative sources of resistance to *Pyricularia oryzae*. *Rice Science*, 29(6), 535-544.

- Tianshun, Z. H. O. U., Dong, Y. U., Ling, L. I. U., Ning OUYANG, G. Y., Meijuan, D. U. A. N., & Dingyang, Y. U. A. N. (2021). CRISPR/Cas9-mediated Editing of AFP1 Improves Rice Stress Tolerance. *Chinese Journal OF Rice Science*, 35(1), 11.
- Tripathi, A. K., Pareek, A., Sopory, S. K., & Singla-Pareek, S. L. (2012). Narrowing down the targets for yield improvement in rice under normal and abiotic stress conditions via expression profiling of yield-related genes. *Rice*, 5, 1-12.
- Wang, S., Li, S., Liu, Q., Wu, K., Zhang, J., Wang, S., ... & Fu, X. (2015). The OsSPL16-GW7 regulatory module determines grain shape and simultaneously improves rice yield and grain quality. *Nature genetics*, 47(8), 949-954.
- Wang, W., Wang, W., Pan, Y., Tan, C., Li, H., Chen, Y., ... & Ma, C. (2022). A new gain-of-function OsGS2/GRF4 allele generated by CRISPR/Cas9 genome editing increases rice grain size and yield. *The Crop Journal*, 10(4), 1207-1212.
- Xing, Y. & Zhang, Q. (2010). Genetic and molecular bases of rice yield. *Annu. Rev. Plant. Biol.*, 61, 421–442.
- Yue, E., Cao, H., & Liu, B. (2020). OsmiR535, a potential genetic editing target for drought and salinity stress tolerance in *Oryza sativa*. *Plants*, 9(10), 1337.
- Zafar, K., Khan, M. Z., Amin, I., Mukhtar, Z., Yasmin, S., Arif, M., ... & Mansoor, S. (2020). Precise CRISPR-Cas9 mediated genome editing in super basmati rice for resistance against bacterial blight by targeting the major susceptibility gene. *Frontiers in plant science*, 11, 575.
- Zafar, K., Sedeek, K. E., Rao, G. S., Khan, M. Z., Amin, I., Kamel, R., ... & Mahfouz, M. M. (2020). Genome editing technologies for rice improvement: progress, prospects, and safety concerns. *Frontiers in Genome Editing*, 2, 5.
- Zeng, X., Luo, Y., Vu, N. T. Q., Shen, S., Xia, K., & Zhang, M. (2020). CRISPR/Cas9-mediated mutation of OsSWEET14 in rice cv. Zhonghua11 confers resistance to *Xanthomonas oryzae* pv. *oryzae* without yield penalty. *BMC Plant Biology*, 20(1), 1-11.
- Zeng, Y., Wen, J., Zhao, W., Wang, Q., & Huang, W. (2020). Rational improvement of rice yield and cold tolerance by editing the three genes OsPIN5b, GS3, and OsMYB30 with the CRISPR–Cas9 system. *Frontiers in plant science*, 10, 1663.
- Zhang, A., Liu, Y., Wang, F., Li, T., Chen, Z., Kong, D., ... & Luo, L. (2019). Enhanced rice salinity tolerance via CRISPR/Cas9-targeted mutagenesis of the OsRR22 gene. *Molecular breeding*, 39, 1-10.
- Zhang, X. Q., Hou, P., Zhu, H. T., Li, G. D., Liu, X. G., & Xie, X. M. (2013). Knockout of the VPS22 component of the ESCRT-II complex in rice (*Oryza sativa* L.) causes chalky endosperm and early seedling lethality. *Molecular biology reports*, 40, 3475-3481.
- Zhou, Y., Xu, S., Jiang, N., Zhao, X., Bai, Z., Liu, J., ... & Yang, Y. (2022). Engineering of rice varieties with enhanced resistances to both blast and bacterial blight diseases via CRISPR/Cas9. *Plant Biotechnology Journal*, 20(5), 876-88.