

Correlation and path coefficient analysis in wheat (*Triticum aestivum* L.) under heat stress condition

Mukti Ram Poudel*, Neeru Lamsal, Anushka Rijal, Prakriti Ghimire, Manisha Ghimire, Kriti Kafle, Karuna Lamsal, Pooja Bhusal

Department of Plant Breeding, Institute of Agriculture and Animal, Science (IAAS), Tribhuvan University, Nepal.

*Correspondence Mukti Ram Poudel muktipoudel8@gmail.com/mukti@iaas.edu.np

Volume: 2, Issue: 2, Pages: 9-14 DOI: https://doi.org/10.37446/corbio/rsa/2.2.2024.9-14 Received: 02 January 2024 / Accepted: 20 March 2024 / Published: 30 June 2024

Background: Twenty distinct wheat genotypes, including three check varieties, were evaluated under late-sown heat stress conditions using correlation and path coefficient analysis to assess the direct and indirect effects of yield-attributing traits on grain yield. The study was conducted during the winter season of 2023 at IAAS Paklihawa Campus, Bhairahawa, Nepal.

Methods: The experiment employed an alpha lattice design with two replications. Heat stress was induced by delayed sowing. Ten randomly selected plants from each genotype were assessed for six traits: plant height, spike length, spikelets per spike, spike weight, thousand grain weight, and tillers per meter square. Correlation analysis and path coefficient analysis were used to determine trait associations with grain yield.

Results: Spikelets per spike was highly positively and correlated with spike weight. Spike weight was highly positively and significantly correlated with thousand grain weight. Besides, grain yield (YLD) showed a significant positive association with tillers per meter square. The other characters such as, spike length, plant height, spike weight, spikelets per spike and thousand grain weight are positively associated with the yield trait.

Conclusion: Breeding efforts should be encouraged for the improvement of positively associated traits with the yield trait in order to boost wheat yields under situations of heat stress.

Keywords: correlation, path coefficient, breeding, genotypes, wheat

Introduction

Wheat is considered as staple food in more than 40 countries of the world which provides basic calories and protein for 85% and 82% of the world population respectively. The annual cereal production has to grow by almost one billion to feed the projected population of 9.1 billion by 2050. Annual wheat production is 781.31 million metric tons all over the world. Wheat is cultivated on 78,000 hectares of land in Nepal with a total production of 218, 500 metric tons (Bhandari & Poudel 2024). In the past 10 years, Nepalese agriculture has seen a decrease in the cropping area of wheat and a very low increase in productivity. MoALD (2020) reports a decrease of 4% cropping area from 731131 ha in 2009/10 to 703992 ha in 2018/19. Similarly, in these ten years, the productivity increase has been very slow at the rate of just 0.102 tons/ha (Poudel et al., 2023). The productivity at the start of the decade (2009/10) was 2.13 tons/ha and reached 2.85 tons/ha by the end of the decade (2018/19) (Bishwas et al., 2021). The area and production of wheat in Nepal have expanded significantly since the introduction of semi-dwarf varieties from Mexico, and it now significantly contributes to the country's food supply, (Pandey et al., 2021). Wheat is composed of Carbohydrates (78%), fat (2%), protein (14%), minerals, B-group vitamins, dietary fiber, riboflavin, thiamine and niacin. Wheat flour is used to prepare bread, biscuits, confectionary products, noodles and vital wheat gluten, also known as seitan (Dhakal et al. 2023). Wheat is known for antinutrients, such as phytic acid, which reduce the absorption of micronutrients, especially

Zn from cereal. The most common deficiencies in the human body are Fe and Zn. In addition, micronutrients such as Zn and Fe accumulate mainly in the germ and bran of the wheat grain, i.e., in the parts that are removed during the processing of the grain, suggesting that whole grain products are an important source of daily intake of these minerals in the human diet. Although the environmental conditions can affect the nutritional composition of wheat grains with its essential coating of bran, vitamins and minerals; it is an excellent health-building food (Petrović et al., 2024). Furthermore, the grain along with stalk and chaff serves as an important raw material for Nepalese industries, and stalk and chaff are also used in the form of mulch, construction material, and animal bedding (Poudel et al., 2023). Wheat grows at temperatures between 15°C and 22°C and it usually grows well in cold, dry conditions. Wheat is cultivated in tropical and subtropical regions that face a variety of abiotic challenges throughout the world. Adverse environmental stress significantly may cause detrimental effects to wheat. The major abiotic stresses include heat, drought, salinity cold, chemical and water excess. However, heat stress is the main abiotic stress affecting wheat production worldwide. Moreover, late sowing of wheat is practiced in Nepal which causes terminal heat stress at the reproductive and ripening stage of wheat (Aryal et al., 2021). The mean ambient temperature is predicted to increase by $1-6^{\circ}C$ by the end of the 21st century. The grain yield of wheat reduces by 6%-10% by just a 1° rise in temperature whereas, in Nepal, a 1-3°C rise in maximum temperature (during heat stress) causes an 8%-31% decrease in wheat yield (Pokhrel et al., 2019) (Bishwas et al., 2021). As a consequence of the low productivity, Nepal has been witnessing a gradual annual increase in wheat imports. The present study aims to understand the correlation and path coefficient analysis to ascertain yield components, as it evaluates the direct and indirect contributions of independent factors to the dependent variable (Puri et al., 2020). Correlation analysis measures would help in the selection of high yielding genotypes by the identification of the association between the grain yields of wheat to its attributing parameters. Path analysis helps to know the direct and indirect (positive or negative) effects of yield-attributing parameters on the grain yield of wheat. It is useful to use genotypic and phenotypic correlations to show how different morpho-physiological traits are related to economic production.

The weather of the Terai region in Nepal indicates that the winter is getting delayed and shorter while warm summer days are becoming longer. This change could be unfavorable to wheat cultivation in the whole Indo Gangetic region including Nepal. Long duration wheat varieties are already suffering due to early heat during the grain filling period (Pokhrel et al., 2019). Late planting of wheat especially in the Terai of Nepal is very common due to several factors such as excess soil moisture just after rice harvesting, delay in rice harvesting due to late maturing varieties or shortage of labours and rice harvesters during rice harvesting season. Wheat crop growth is highly dependent on temperature regimes and high temperature reduces the physiological maturity days as well as the grain filling duration, as a result reducing wheat yield (Poudel et al., 2024). Under such situations, wheat varieties with shorter maturity or a faster grain filling rate would be desirable. Therefore, breeding for high-temperature tolerant wheat genotypes having disease resistance and stay green traits that are capable of producing enough endosperm, bold grains within short grain filling duration was conducted to identify terminal heat tolerance in breeding lines of wheat.

Materials and methods

The experiment was carried out on the research field at the Paklihawa Campus, Institute of Agriculture and Animal Science (IAAS), Rupandehi, The experiment site is located at 104 meters above sea level and at 27 29'02" N and 83 27' 17" E. A set of twenty wheat genotypes (Table 1) was obtained from the National Wheat Research Program (NWRP), Bhairahawa, Nepal including RR, Bhrikuti and Gautam as check varieties

S.No.	Genotypes	S.No.	Genotypes
1	BL_5106	11	NL_1447
2	BL 5099	12	NL ¹⁴⁴⁵
3	BL_4984	13	NL_1506
4	Bhrikuti	14	NL_1504
5	NL 1437	15	NL 1503
6	NL ¹⁴⁰²	16	NL 1501
7	Gautam	17	RR_21
8	BL_5116	18	NL_1512
9	NL_1492	19	NL_1509
10	NL_1488	20	NL_1508

Table1. List of <u>genotype (Source:</u>	NWRP, Bhairahawa	a) used in experiment
--	------------------	-----------------------

The experiment was conducted on alpha lattice design (Table 2) with two replications of a plot size 5m² with a dimension of 2.5 x 2m. The inter replication spacing was maintained at 1 m while interplot spacing was 25cm. Heat stress condition was created by sowing wheat genotypes one month later (26th December) than the normal condition (25th November) so as to coincide with heat waves with reproductive stage of the wheat crop. Ten randomly selected plants were tagged independently for each genotype and yield-attributing traits were documented including Plant height (Ph), Spike length (SI), Spike Weight (SW), Thousand Grain Weight (TGW), Number of Spikes Per Meter Square (NSPMS) and Grain Yield (GY).

Table 2. Layout of Experimental site							
Replication 1]	Replication 2			
T1	T2	T3	T4	T1	T2	T3	T4
T8	Τ7	T6	T5	T8	T7	T6	T5
Т9	T10	T11	T12	Т9	T10	T11	T12
T16	T15	T14	T13	T16	T15	T14	T13
T17	T18	T19	T20	T17	T18	T19	T20

The recommended dose of fertilizer was applied in each individual plot with a dosage of 100:50:25 kgNPK/ha,and compost manure was applied at a rate of 5 tons per hectare. Prior to seeding, all of the phosphorus, potash, and half of the nitrogen dose were administered. Two divided doses of the remaining nitrogen were administered: the first quarter at 30 DAS and the last dosage at 70 DAS. Irrigation was provided as per the recommendation of the National Wheat Research Program (NWRP) MS Excel 2016 was used for data entry and processing. Using SPSS Version 20, the association between the yield and yield-attributing characteristics was determined. The path analysis was done using Microsoft Excel 2016.

Results

The yield-contributing morphological features are essential for high yielding variety breeding to be successful. The quantity of tillers/m², the number of grains per spike, the weight of 1000 grains, plant height, and spike length are some of the factors that influence the complicated characteristics of wheat grain production. The correlation analysis (Table 3) shows a positive association of grain yield with plant height (0.160).

Table 5. Correlation between different quantitative traits							
	Ph	SL	SPS	SW	TGW	TPM	YLD
Ph	1	0.134	-0.020	-0.073	-0.020	-0.052	0.160
SL	0.134	1	0.187	0.237	0.023	-0.076	0.199
SPS	-0.020	0.187	1	0.538**	0.042	-0.168	0.073
SW	-0.073	0.237	0.538**	1	0.495**	-0.188	0.150
TGW	-0.020	-0.023	0.042	0.495**	1	-0.215	0.060
TPM	-0.052	-0.076	-0.168	-0.188	-0.215	1	0.336*
YLD	0.160	0.199	0.073	0.150	0.060	0.336*	1

Table 3. Correlation between different quantitative traits

Where, Ph = Plant Height, SL = Spike Length, SPS = Spikelets per Spike, SW = Spike Weight, TGW = Thousand Grain Weight, TPM = Tillers Per Meter, YLD = Yield

Plant height showed a weak positive correlation with yield (Table 3) where Plant height was negatively correlated with spikelet per spike, spike weight, thousand grain weight, and tillers per metre square. Spike length was positively correlated with plant height, spikelet per spike, spike weight, thousand grain weight and yield where spike length was negatively correlated with tillers per square metre. Spikelet per spike was highly positively significantly correlated with spike weight (0.538**) and positively with spike length, thousand grain weight and yield where spikelet per spike was negatively correlated with plant height and tillers per metre.

Spike Weight was highly positively significantly correlated with Thousand grain weight (0.495**) and positively correlated with Spike length and yield where negatively correlated with plant height and tillers per meter square. Thousand grain weight showed a positive significant correlation with spike weight, and a weak negative correlation with plant height and tillers per meter square. Tillers per meter square is positively significantly correlated with yield (0.336*).

Path coefficient analysis aims to find out what affects the endogenous variables, that is, how the exogenous variables work together (their covariance) and which paths are important (determined in part by the variances of the exogenous variables). The analysis (Table 4, Figure 1) revealed that tillers per meter exhibited the highest positive direct effect on

grain yield (0.4030). Following TPM, plant height showed a positive direct effect (0.1857). Spike weight also contributed positively and directly to grain yield (0.1707). Other traits with positive direct effects on grain yield included spike length (0.0758), thousand grain weight (0.0629), and spikelets per spike (0.0358).

Table 4. I all analysis of unferent quantitative trait on grain yield							
Trait	Ph	SL	SPS	SW	TGW	TPM	
ViaPh	0.1857	0.0248	-0.0037	-0.0135	-0.0037	-0.0097	
ViaSL	0.0101	0.0758	0.0142	0.0180	0.0018	-0.0057	
ViaSPS	-0.0007	0.0067	0.0358	0.0192	0.0015	-0.0060	
ViaSW	-0.0124	0.0405	0.0918	0.1707	0.0845	-0.0320	
ViaTGW	-0.0012	0.0015	0.0026	0.0311	0.0629	-0.0135	
ViaTPM	-0.0211	-0.0305	-0.0677	-0.0756	-0.0866	0.4030	
ViaYLD	0.1604	0.1187	0.0729	0.1500	0.0604	0.3360	

Table 4. Path analysis of different quantitative trait on grain yield

Where, Ph = Plant Height, SL = Spike Length, SPS = Spikelets Per Spike, SW = Spike Weight, TGW = Total Grain Weight, TPM = Tillers Per Meter, YLD = Yield. Residual effect=0.3754



Figure 1. Graphical representation of path coefficient analysis

Plant height exerted negative indirect effects on grain yield via spikelets per spike (-0.0037), spike weight (-0.0135), thousand grain weight (-0.0037), and tillers per meter (-0.0097). its positive indirect effect was observed via spike length (0.0248). Spike length showed a positive indirect effect on grain yield via spikelets per spike (0.0142) and spike weight (0.0180). Conversely, it exhibited a negative indirect effect via tillers per meter (-0.0057). Spikelets per spike had positive indirect effects on grain yield through spike weight (0.0192) and thousand grain weight (0.0015). However, it showed negative indirect effects via plant height (-0.0007) and tillers per meter (-0.0060). spike weight (0.0845). It had a negative indirect effect via tillers per meter (-0.0320). Thousand grain weight showed a positive indirect effect via tillers per meter (-0.0135). Tillers per meter exhibited significant negative indirect effects on grain yield via spikelets per spike (-0.0135). Tillers per meter exhibited significant negative indirect effects on grain yield via spikelets per spike indirect effects on grain yield via spikelets per spike indirect effects on grain yield via spikelets per spike (-0.00135). Tillers per meter (-0.0756), and thousand grain weight (-0.0866).

Discussion

The correlation coefficient measures the strength and direction of a linear relationship between two variables. Correlation refers to the fact that there is a link between various events. One of the tools to infer whether such a link exists is correlation analysis. Practical simplicity is undoubtedly one of its main advantages. Plant height was highly positively correlated with grain yield as reported by Mukherjee et al. (2008). The taller the plant, the higher will be the photosynthate area due to the long stem. According to Poudel et al. (2023), SL was positively associated with grain yield which is due to the accumulation of more grains in long spikes. In contrast, a shorter spike accumulates fewer grains and leads to a lower yield of wheat. Path analysis is a statistical technique for examining and testing relationships among a set of observed variables. Relationships or paths may be described as representing direct or

indirect associations. An indirect relationship occurs when a variable is linked to another variable via a third variable, which in turn is directly associated with the outcome variable. The sum of all direct and indirect relationships, in turn, yields a variable's total association. Because path analysis allows the study of multiple direct and indirect relationships between variables simultaneously, it is regarded as a more complex technique than typical regression analysis. Perhaps the greatest advantage of path analysis over regression analysis is its superior ability to detect associations. With regression, the absence of a direct correlation of one variable with another is usually interpreted as a null association (Wooldredge, 2021). Thousand grain weight had a direct positive effect on grain yield (0.0629) but Tillers per meter square showed an indirect negative effect on plant height and tillers per meter square. According to Guendouz et al. (2013), the direct effect of 1000 grain weight upon grain yield was high and positive. Tillers per meter square had a highest direct effect on grain yield (0.4030). It may be concluded from the present studies that plant height, peduncle length, tillers per plant, spike length, spike weight and number of grains per spike appeared to contribute to the grain yield. Therefore indirect selection for higher grain yield may be effective for improving these characters, as shown by Ozukum et al., 2019.

Conclusion

Wheat (*Triticum aestivum*) is one of the most important staple food crops in the lowland Terai area of Nepal. However, the nation's production has failed to meet the growing demand. From a South Asian regional perspective, the average production is low with significant spatial and temporal variability. Heat stress condition provides the adaptability and tolerance of wheat in a heat stress condition. Path coefficient analysis also evaluated that thousand grain weights were highly positively correlated with spike weight but negatively correlated with plant height and tillers per meters square. Tillers per meter square is positively correlated with yield but negatively correlated with plant height, spike length, spikelet per spike, spike weight and tillers per meter square. Plants with a tall height should be promoted to obtain maximum higher yield.

Acknowledgment

The authors acknowledge the National Wheat Research Program (NWRP) for providing genetic materials, the Nepal Academy of Science and Technology (NAST) for financial support to conduct research and the Institute of Agriculture and Animals Science (IAAS), Paklihawa campus for providing field for experiment.

Author contributions

All authors have contributed equally to the present research work. The final manuscript was prepared by Pooja Bhushal and MR Poudel.

Conflict of interest

The authors declare no conflict of interest.

Ethics approval

Not applicable.

AI tool usage declaration

No AI tools have been used in manuscript preparation.

References

Aryal, S., Dhungel, B., Subedi, K. R., Lamichhane, P., & Bhattarai, S. (2021). Response of Wheat Parameters To Sowing Date and Irrigation Supplied in Terai Region of Nepal. *Tropical Agrobiodiversity*, 2(1), 07–09. https://doi.org/10.26480/trab.01.2021.07.09

Bhandari, R., & Poudel, M. R. (2024). Genotype × environment interaction and selection parameters for high yielding wheat genotypes under irrigated and heat stress environment. *Journal of Sustainable Agriculture and Environment*, 3(1). https://doi.org/10.1002/sae2.12098

Bishwas, K. C., Poudel, M. R., & Regmi, D. (2021). AMMI and GGE biplot analysis of yield of different elite wheat line under terminal heat stress and irrigated environments. *Heliyon*, 7(6), e07206. https://doi.org/10.1016/j.heliyon.2021.e07206

Dhakal, A., Bhandari, R., Nyaupane, S., Paudel, H., Panthi, B., & Poudel, M. R. (2023). Agro Morphological Analysis of Wheat Genotype Under Heat Stress Condition. *Journal of Food and Agriculture Research*, *3*(1), 35–43.

Guendouz, A., Djoudi, M., Guessoum, S., Maamri, K., Fellahi, Z., Hannachi, A., & Hafsi, M. (2013). Durum Wheat (*Triticum durum* Desf.) Evaluation under Semi Arid Conditions in Eastern Algeria by Path Analysis. *Journal of Agriculture and Sustainability*, 3(2).

Kumar, A., Yadav, V. K., Maurya, C. L., Singh, S. V., Shweta, Kumar, J., & Naik, B. S. D. (2024). Estimate of Genotypic and Phenotypic Correlation and Path Coefficients in Bread Wheat (*Triticum aestivum* L.). *Journal of Advances in Biology & Biotechnology*, *27*(3), 149–170. https://doi.org/10.9734/jabb/2024/v27i3730

MoALD, 2020. Area, Production and Yield by Major Cereal Crops, Last Ten Years. https://s3-ap-southeast-1.amazonaws.com/prod-gov-agriculture/server-assets/publicat ion-1568704754479-585b7.xlsx

Mukherjee, S., Suparna Gupta, S. G., & Niladri Bhowmik, N. B. (2008). Genetic variability of important quantitative characters in modern wheat (*Triticum aestivum* L.) cultivars in Tarai Region of West Bengal. *Environment and Ecology*, 26 (2), 683-686.

Ozukum, W., Avinashe, H., Dubey, N., Kalubarme, S., & Kumar, M. (2019). Correlation and path coefficient analyses in bread wheat (*Triticum aestivum* L.). *Plant Archives*, *19*(2), 3033–3038.

Pandey, D., Pant, K. R., Bastola, B. R., Giri, R., Bohara, S., Shrestha, S., Hamal, G. B., & Shrestha, J. (2021). Evaluation of bread wheat genotypes under rain-fed conditions in Terai districts of Nepal. *Journal of Agriculture and Natural Resources*, *4*(2), 303–315. https://doi.org/10.3126/janr.v4i2.33946

Petrović, S., Vila, S., Grubišić Šestanj, S., & Rebekić, A. (2024). Variation in Nutritional Value of Diverse Wheat Genotypes. *Agronomy*, 14(2). https://doi.org/10.3390/agronomy14020311

Pokhrel, D., Raj Pant, K., Raman Upadhyay, S., Pandey, D., Raj Gautam, N., Khatri, N., Prasad Yadav, R., Ram Ghimire, B., Prasad Poudel, R., Prasad Poudel, B., Prasad Paudel, G., Raj Yadav, D., & Basnet, R. (2019). *Development of suitable wheat varieties for terminal heat stress environment in Terai region of Nepal. January 2022*, 20–21.

Poudel, M.R., Neupane M.P., Panthi B., Bhandari R.K., Nyaupane S.L., Dhakal A., & Paudel H.K.. (2023). "Identification of Drought Tolerant Wheat (*Triticum Aestivum*) Genotypes Using Stress Tolerance Indices in the Western Terai Region of Nepal." *Research on Crops* 24(4), 652–59.

Poudel, M. R., Poudel, H. K., & Bhandari, R. (2024). Impact of terminal heat stress on performance of Nepalese wheat (Triticum aestivum L.) genotypes. *Research on Crops*, 25(1), 1-11.

Poudel, M. R., Neupane, M. P., Paudel, H., Bhandari, R., Nyaupane, S., Dhakal, A., & Panthi, B. (2023). Agromorphological analysis of wheat (*Triticum aestivum* L.) genotypes under combined heat and drought stress conditions. *Farming & Management*, 8(2), 72-80.

Puri, R. R., Tripathi, S., Bhattarai, R., Dangi, S. R., & Pandey, D. (2020). Wheat Variety Improvement for Climate Resilience. *Asian Journal of Research in Agriculture and Forestry*, *6*(2), 21–27. https://doi.org/10.9734/ajraf/2020/v6i230101

Wooldredge, J. (2021). Path Analysis. *The Encyclopedia of Research Methods in Criminology and Criminal Justice: Volume II: Parts 5-8, November 2017*, 515–522. https://doi.org/10.1002/9781119111931.ch105