



## Effect of different sources of planting material on growth, yield and quality attributes in cassava (*Manihot esculenta* Crantz.)

Sundaramoorthy Nanthakumar<sup>1</sup>, Krithika K<sup>2</sup>, Muthusamy Prabhu<sup>3\*</sup>, Govindasamy Ashok Kumar<sup>4</sup>

<sup>1</sup>Professor (Horticulture), TNAU - Agricultural Research Station, Virinjipuram, Vellore District, Tamil Nadu, India.

<sup>2</sup>Post Graduate Student (Vegetable Science), TNAU, Coimbatore, Tamil Nadu, India.

<sup>3</sup>Associate Professor (Horticulture), TNAU - Horticultural College and Research Institute, Coimbatore, Tamil Nadu, India.

<sup>4</sup>Assistant Professor (Horticulture), TNAU - Horticultural College and Research Institute, Coimbatore, Tamil Nadu, India.

**\*Correspondence**

Muthusamy Prabhu  
prabhu.m@tnau.ac.in

Volume: 11, Issue: 3, Pages: 10-17

DOI: <https://doi.org/10.37446/jinagri/rsa/11.3.2024.10-17>

Received: 25 March 2024 / Accepted: 15 August 2024 / Published: 30 September 2024

**Background:** Cassava is an industrial tropical tuber crop and it is used for human and animal consumption: but also used as a raw material for various agro-based industries. Cassava is commercially propagated by using stem cuttings. These stem cuttings are often infected with viral and bacterial diseases which affect yield adversely. With this background, the present experiment was taken up to identify an elite virus free planting material with improved yield and qualitative characters in two promising cultivars of tapioca *viz.*, H 226 and CO 2 using tissue cultured plants and different types of setts from field grown plants as planting materials.

**Methods:** An experiment was carried out to assess the effect of planting materials on the growth, yield and quality of Cassava varieties *viz.*, H 226 and CO 2. The experiment was laid out in a Factorial Randomized Block Design with four treatments in two varieties and three replications. Observations were recorded on various growth, yield and quality attributes of the crop.

**Results:** Significant results on plant height, stem girth, number of branches and number of leaves are recorded with tissue cultured plants. There is no significant difference among the treatments for starch, HCN and protein contents. There is a significant difference in starch content among the varieties. The variety CO 2 possessed more starch content than H 226.

**Conclusion:** Among the four treatments (different kinds of planting materials) followed in this study, the treatment T<sub>1</sub> Tissue cultured plants recorded lesser incidence of cassava mosaic virus, more number of fibrous roots, highest values for plant height, stem girth and tuber yield. Among the varieties, CO 2 showed the least incidence of cassava mosaic virus. The CO 2 tissue culture plants could give better yield in the field conditions.

**Keywords:** cassava, tissue culture, mosaic disease, yield

### Introduction

Cassava (*Manihot esculenta* Crantz.) is an industrial tropical tuber crop and it is used for human and animal consumption: but also used as a raw material for various agro-based industries. The percentage of carbohydrate production from the unit area of cassava is 40 per cent higher than rice and 25 per cent more than maize (Tonukari, 2004). Cassava tubers are the cheapest source of calories (Tonukari, 2004). Cassava, the third main source of carbohydrates in Africa, provides daily nutrition for over 700 million people on the continent and a vital source of

income for subsistence farmers (Szyniszewska, 2020). Africa is a major contributor of cassava tuber production (57%), followed by Asia (25%). Tapioca is being cultivated in 15.70 million hectares of area and production was 158 million tonnes with an average productivity of 10t/ha in the world. In India, presently, cassava is cultivated in an area of 0.163 m ha, with a production of 4.98 m tons and productivity of 30.55 t ha<sup>-1</sup> (Radhakrishnan et al., 2022). Major cassava production states in India are Kerala, Tamil Nadu and Andhra Pradesh. True seeds of cassava are not used for commercial cultivation due to high seed dormancy and very poor germination. Commercially, farmers are practicing 8-10 cm long cuttings from the mature part of stem and planted in vertical position in specially prepared nursery bed at closer spacing of 5 cm. The rooted cuttings cannot be transported from one place to other place because of bare roots. Cassava is highly susceptible to cassava mosaic disease (CMD) and disease spreads through infected setts and subsequently transmitted by white flies (Pugalendhi & Velmurugan, 2020). In many the cases tissue culture techniques are used for large scale propagation of vegetatively propagated crops (Faria & Illg, 1995), to eliminate the virus and other pathogens. With this background, the present experiment was taken up to identify an elite virus free planting material with improved yield and qualitative characters in two promising cultivars of tapioca viz., H 226 and CO 2 using tissue cultured plants and different types of setts from field grown plants as planting materials.

## Materials and Methods

The present experiment was carried out at College Orchard of Tamil Nadu Agricultural University, Coimbatore. The maximum temperature of the location fluctuated between 25.5°C and 34.4°C, with a minimum temperature ranging between 16.5°C and 23.4°C. Relative humidity ranged from 30 to 93 per cent. The soil of the experimental field was clay loam in texture and the pH ranged 8.0- 8.5. The experiment was laid out in a factorial randomized block design with a plot size of 4m x 4m and replicated thrice. Following promising varieties of cassava were taken for the experiment.

V<sub>1</sub> -H 226 released from ICAR Central Tuber Crop Research Institute, Thiruvananthapuram  
 V<sub>2</sub> - CO 2 released from Tamil Nadu Agricultural University, Coimbatore

The planting materials taken for the study are as follows.

T<sub>1</sub> - Tissue cultured plants  
 T<sub>2</sub> . Two budded setts from tissue cultured grown plants  
 T<sub>3</sub> . Six budded setts from normal field grown plants and  
 T<sub>4</sub>-Two budded setts from normally field grown plants.

The setts with different nodes were planted, irrigated daily and kept under shade net for 25 days in the nursery and then they were transplanted the main field. In the case of tissue culture plants were kept under shade net for secondary hardening. Two months old plants were planted in the main field. Well decomposed farm yard manure was applied at the rate of 25t/ha as a basal dose before final ploughing and basal dose of 45:80:120 kg/ha N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O was applied. Then the tissue cultured plants and setts were planted at a spacing of 90 x 75 cm at the ridges. The growth characters viz., plant height, stem girth, number of branches, and number of leaves were recorded on 270<sup>th</sup> day after transplanting. Cassava mosaic disease incidence was counted from infected plants in a plot. The cassava mosaic incidence was calculated by the formula 'CMD incidence (%) =(No of plants infected/ Total no. of plants plot<sup>-1</sup>) x 100'.

The yield contributing parameters viz., tuber yield per plot, number of tubers per plant, tuber yield per plant, length and girth of tuber are recorded during harvest. The harvesting was done on fifth April, 2022. Similarly, the quality attributing traits viz., starch content (Hernández-Carmona et al., 2017), protein content of tubers were estimated as per standard procedures (Lowry et al., 1951). The HCN content in the flesh of the tubers was estimated at harvesting stage, from the tuber the samples were taken and by the method described by Indira & Sinha (1969). The benefit cost ratio was worked out in terms of total expenditure. The observations recorded on various attributes were statistically analysed by the method recommended by Panse & Sukhatme (1954).

## Results

The treatment T<sub>1</sub> (tissue cultured plants) has recorded maximum plant height, stem girth, number of branches and number of leaves. Among the two varieties, H 226 (V<sub>1</sub>) has recorded the highest plant height and the variety CO 2 (V<sub>2</sub>) has recorded the higher stem girth, number of branches and number of leaves. The interaction between treatment and varieties were significant. Tissue cultured plants (T<sub>1</sub>) of cv. H 226 (V<sub>1</sub>) recorded the highest plant height and tissue cultured plants of variety CO 2 recorded higher stem girth, number of branches and number of leaves (Table 1,2,3,4).

**Table 1. Effect of planting materials on plant height (cm) in cassava**

Treatments	90 DAT			180 DAT			270 DAT		
	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean
T <sub>1</sub>	62	55	57	143	127	135	215	199	207
T <sub>2</sub>	40	47	44	124	118	121	193	191	192
T <sub>3</sub>	59	40	51	132	125	128	205	198	202
T <sub>4</sub>	39	32	36	126	118	122	193	192	192
Mean	50	44	47	131	122	127	202	195	198
	V	T	VxT	V	T	VxT	V	T	VxT
S.E(d)	4.6	6.5	9.2	0.6	0.8	1.2	1.2	1.7	2.4
<b>CD (5%)</b>	<b>9.9</b>	<b>14.0</b>	<b>19.8</b>	<b>1.3</b>	<b>1.8</b>	<b>2.5</b>	<b>2.6</b>	<b>3.7</b>	<b>5.3</b>

**Table 2. Effect of planting materials on stem girth (cm) in cassava treatments**

	90DAT			180DAT			270DAT		
	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean
T <sub>1</sub>	3.9	4.7	4.3	9.0	9.7	9.4	11.1	13.2	12.1
T <sub>2</sub>	3.9	3.8	3.8	7.5	7.1	7.3	9.4	11.0	10.2
T <sub>3</sub>	3.8	4.1	3.9	8.0	8.3	8.1	10.4	11.7	11.1
T <sub>4</sub>	3.5	3.7	3.6	7.3	7.1	7.2	9.1	10.9	10.0
Mean	3.8	4.0	4.9	8.0	8.0	8.0	8.0	9.4	8.7
	V	T	VxT	V	T	VxT	V	T	VxT
S.Ed	0.18	0.26	0.36	0.07	0.10	0.15	0.17	0.24	0.35
<b>CD (5%)</b>	<b>0.39</b>	<b>0.55</b>	<b>0.79</b>	<b>0.16</b>	<b>0.23</b>	<b>0.32</b>	<b>0.37</b>	<b>0.53</b>	<b>0.75</b>

**Table 3. Effect of planting materials on the total number of branches in cassava**

Treatments	90DAT			180DAT			270DAT		
	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean
T <sub>1</sub>	2.80	3.26	3.03	3.26	4.43	3.84	7.16	9.96	8.56
T <sub>2</sub>	1.60	2.90	2.25	2.56	3.83	3.19	5.76	8.46	7.11
T <sub>3</sub>	2.30	3.23	2.76	3.16	4.40	3.78	6.73	9.96	8.34
T <sub>4</sub>	1.50	2.10	1.80	2.46	3.50	2.98	5.70	8.40	7.05
Mean	2.05	2.87	2.46	2.86	4.04	3.45	6.33	9.19	7.76
	V	T	VxT	V	T	VxT	V	T	VxT
S.Ed	0.15	0.22	0.31	0.10	0.15	0.21	0.14	0.21	0.29
<b>CD (5%)</b>	<b>0.33</b>	<b>0.47</b>	<b>0.67</b>	<b>0.22</b>	<b>0.32</b>	<b>0.45</b>	<b>0.31</b>	<b>0.45</b>	<b>0.63</b>

**Table 4. Effect of planting materials on the number of leaves per plant in cassava**

Treatments	90DAT			180DAT			270DAT		
	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean
T <sub>1</sub>	36.0	42.1	39.0	102.0	125.2	113.6	69.2	70.2	69.7
T <sub>2</sub>	29.3	31.0	30.1	87.9	109.0	98.4	45.8	59.4	52.6
T <sub>3</sub>	34.1	39.3	36.7	99.2	119.3	109.3	59.3	69.5	64.4
T <sub>4</sub>	28.1	29.9	29.0	83.6	99.9	91.7	43.5	58.3	50.9
Mean	31.8	35.6	33.7	93.2	113.4	103.3	54.4	64.4	59.4
	V	T	VxT	V	T	VxT	V	T	VxT
S.Ed	2.17	3.07	4.34	1.56	2.79	3.45	2.1	3.01	4.12
<b>CD (5%)</b>	<b>4.66</b>	<b>6.59</b>	<b>9.33</b>	<b>3.54</b>	<b>5.98</b>	<b>8.72</b>	<b>4.2</b>	<b>5.99</b>	<b>9.10</b>

V<sub>1</sub> - H 226, V<sub>2</sub> - CO 2, DAT- Days after transplantingT<sub>1</sub>- Tissue cultured plantsT<sub>2</sub>- Two budded setts from tissue cultured plantsT<sub>3</sub>- Six budded setts from field grown plantsT<sub>4</sub>- Two budded setts from field grown plants

The treatment T<sub>3</sub> (Six budded setts from field grown plants) recorded the highest CMD incidence and the least CMD incidence was recorded by T<sub>1</sub> (tissue cultured plants). Among the varieties, H 226 (V<sub>1</sub>) recorded the higher CMD incidence and CO 2 (V<sub>2</sub>) recorded the lesser CMD incidence (Table 5).

**Table 5. Effect of planting materials on CMD incidence (%) in cassava**

Treatments	90DAT			180DAT			270DAT		
	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean
T <sub>1</sub>	1.10	0.00	0.55	2.20	0.90	1.55	3.00	1.00	2.00
T <sub>2</sub>	21.33	1.01	11.17	22.43	2.50	12.46	23.30	3.33	13.31
T <sub>3</sub>	45.50	10.45	27.97	55.20	15.50	35.35	78.33	22.33	50.33
T <sub>4</sub>	44.90	10.33	27.61	49.90	15.22	32.56	77.93	22.01	49.97
Mean	18.20	5.44	16.82	33.43	8.53	20.48	45.64	12.16	28.90
	V	T	VxT	V	T	VxT	V	T	VxT
S.Ed	0.19	0.22	0.31	0.44	0.52	0.64	1.01	1.18	1.40
<b>CD (5%)</b>	<b>0.45</b>	<b>0.51</b>	<b>0.63</b>	<b>0.90</b>	<b>1.10</b>	<b>1.21</b>	<b>1.30</b>	<b>1.65</b>	<b>2.01</b>

V<sub>1</sub> - H 226, V<sub>2</sub> - CO 2T<sub>1</sub>- Tissue cultured plantsT<sub>2</sub>- Two budded setts from tissue cultured plantsT<sub>3</sub>- Six budded setts from field grown plantsT<sub>4</sub>- Two budded setts from field grown plants

DAT- Days after transplanting

The treatments T<sub>1</sub> (tissue cultured plants) registered significantly more number of tubers per plant followed by T<sub>3</sub> (Six budded setts from field grown plants). The significant difference was higher in CO 2 (V<sub>2</sub>) which was followed by H 226 (V<sub>1</sub>). As far as the interaction effect, CO 2 (V<sub>2</sub>) along with T<sub>1</sub> (Tissue cultured plants) produced the maximum number of tubers per plant (Table 6).

**Table 6. Effect of planting material on tuber yield of cassava**

Treatments	No. of tubers plant <sup>-1</sup>			Tuber yield plant <sup>-1</sup>			Tuber yield plot <sup>-1</sup> (kg)			Tuber yield (t ha <sup>-1</sup> )		
	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean
T <sub>1</sub>	7.7	10.7	9.2	5.2	6.0	5.6	70.2	87.8	79.0	43.8	54.9	49.3
T <sub>2</sub>	5.5	6.7	6.1	3.8	4.9	4.3	58.8	79.1	68.9	36.7	49.4	43.0
T <sub>3</sub>	7.3	9.7	8.5	4.9	5.5	5.2	67.2	83.1	75.1	42.0	51.9	46.9
T <sub>4</sub>	5.3	6.3	5.8	3.4	4.1	3.7	54.7	74.9	64.8	34.2	46.8	40.5
Mean	6.4	8.3	7.4	3.5	5.1	5.3	62.7	81.2	71.9	39.2	50.7	44.9
	V	T	VxT	V	T	VxT	V	T	VxT	V	T	VxT
S.Ed	0.30	0.42	0.60	0.25	0.34	0.50	0.08	0.17	0.19	0.19	0.32	0.28
<b>CD (5%)</b>	<b>0.64</b>	<b>0.91</b>	<b>1.28</b>	<b>0.52</b>	<b>0.70</b>	<b>0.94</b>	<b>0.20</b>	<b>0.24</b>	<b>0.31</b>	<b>0.35</b>	<b>0.47</b>	<b>0.72</b>

V<sub>1</sub> - H 226, V<sub>2</sub> - CO 2T<sub>1</sub>- Tissue cultured plantsT<sub>2</sub>- Two budded setts from tissue cultured plantsT<sub>3</sub>- Six budded setts from field grown plantsT<sub>4</sub>- Two budded setts from field grown plants

The treatment T<sub>1</sub> (Tissue cultured plants) registered significantly higher tuber yield followed by T<sub>3</sub> (Six budded setts from field grown plants). Among the two varieties, CO 2 (V<sub>2</sub>) gave maximum yield and H 226 recorded lowest value. Within the interaction effect, CO 2 (V<sub>2</sub>) along with T<sub>1</sub> (Tissue cultured plants) gave the maximum tuber yield. The lowest tuber yield was recorded in H 226 (V<sub>1</sub>) along with T<sub>4</sub> (Two budded setts from field grown plants).

**Table 7. Effect of planting material on tuber characters of cassava**

Treatments	Tuber length (cm)			Tuber girth (cm)		
	V <sub>1</sub>	V <sub>2</sub>	Mean	V <sub>1</sub>	V <sub>2</sub>	Mean
T <sub>1</sub>	38.3	35.4	36.9	17.6	20.2	18.9
T <sub>2</sub>	36.1	34.0	35.1	16.3	18.2	17.3
T <sub>3</sub>	37.1	34.2	35.7	17.1	19.4	18.2
T <sub>4</sub>	34.4	33.2	33.8	15.4	17.4	16.4
Mean	36.5	34.2	35.3	16.6	18.8	17.7
	V	T	VxT	V	T	VxT
S.Ed	0.19	0.27	0.39	0.18	0.26	0.36
<b>CD (5%)</b>	<b>0.42</b>	<b>0.59</b>	<b>0.84</b>	<b>0.39</b>	<b>0.55</b>	<b>0.79</b>

Among the treatments,  $T_1$  (Tissue cultured plants) recorded the maximum tuber length and tuber girth followed by  $T_3$  (Six budded setts from field grown plants) than the other two treatments.  $H\ 226 (V_1)$  recorded the maximum tuber length followed by  $CO\ 2 (V_2)$ . The interaction between treatments and cultivars was statistically significant. The maximum tuber length was recorded in  $T_1$  (Tissue cultured plants) along with  $H\ 226 (V_1)$ . The shortest tuber length was recorded in  $CO\ 2 (V_2)$  combined with  $T_4$  (Two budded setts from field grown plants). The higher tuber girth was recorded in  $CO\ 2 (V_2)$  which was followed by  $H\ 226 (V_1)$ . The variety  $CO\ 2 (V_2)$  combined with  $T_1$  (Tissue cultured plants) recorded the highest tuber girth and the least tuber girth was recorded in  $H\ 226 (V_1)$  along with  $T_4$  (Two budded setts from field grown plants) (Table 7).

**Table 8. Effect of planting material on tuber quality characters of cassava**

Treatments	Starch content (%)			Protein content (%)			HCN content ( $\mu\text{g g}^{-1}$ )		
	$V_1$	$V_2$	Mean	$V_1$	$V_2$	Mean	$V_1$	$V_2$	Mean
$T_1$	29.8	34.1	31.9	21.1	26.1	23.6	22.5	10.1	16.3
$T_2$	28.3	33.3	30.8	20.0	26.0	23.0	21.9	10.0	15.9
$T_3$	29.1	34.1	31.6	20.1	25.9	23.0	22.7	10.0	16.4
$T_4$	27.7	32.9	30.3	20.1	25.8	23.0	21.9	10.0	15.9
Mean	28.7	33.6	31.2	20.3	25.9	23.1	22.3	10.0	16.1
	V	T	VxT	V	T	VxT	V	T	VxT
S.Ed	0.20	0.28	0.40	0.04	0.03	0.07	0.12	0.19	0.21
<b>CD (5%)</b>	<b>0.43</b>	<b>NS</b>	<b>0.86</b>	<b>0.08</b>	<b>NS</b>	<b>0.14</b>	<b>0.45</b>	<b>NS</b>	<b>0.63</b>

$V_1$  -  $H\ 226$ ,  $V_2$  -  $CO\ 2$

$T_1$ - Tissue cultured plants

$T_2$ - Two budded setts from tissue cultured plants

$T_3$ - Six budded setts from field grown plants

$T_4$ - Two budded setts from field grown plants

There is no significant difference among the treatments for starch, HCN and protein contents. Among the varieties,  $CO\ 2 (V_2)$  recorded higher starch and protein contents followed by  $H\ 226 (V_1)$ . But there is a significant difference among the varieties,  $H\ 226 (V_1)$  recorded higher HCN content followed by  $CO\ 2 (V_2)$  (Table 8).

**Table 9. Economic of planting material in cassava**

Sl. No.	Parameters	$V_1$				$V_2$			
		$T_1$	$T_2$	$T_3$	$T_4$	$T_1$	$T_2$	$T_3$	$T_4$
1.	Gross cost of production (Rs. $\text{ha}^{-1}$ )	44129	41023	43290	39925	50632	49578	49974	49120
2.	Yield of produce ( $\text{t ha}^{-1}$ )	43.8	36.7	42.0	46.8	54.9	49.4	51.9	46.8
3.	Selling price (Rs. $\text{t}^{-1}$ )	4500	4500	4500	4500	4500	4500	4500	4500
4.	Gross income (Rs. $\text{ha}^{-1}$ )	197100	165150	189000	210600	247050	222300	233550	210600
5.	Gross net income (Rs. $\text{ha}^{-1}$ )	152971	124127	145710	170675	196418	172722	183576	161480
6.	Gross benefit - cost ratio	3.5	3.0	3.4	4.3	3.9	3.5	3.7	3.3

$V_1$  -  $H\ 226$ ,  $V_2$  -  $CO\ 2$

$T_1$ - Tissue cultured plants

$T_2$ - Two budded setts from tissue cultured plants

$T_3$ - Six budded setts from field grown plants

$T_4$ - Two budded setts from field grown plants

The economics worked out for different treatments showed that  $T_1$  (Tissue cultured plants) was found to be economically superior in term of benefit cost ratio followed by  $T_3$  (Six budded setts from field grown plants) (Table 9) in  $CO\ 2 (V_2)$ .

## Discussion

In the present study, there was a difference between the two varieties with respect to growth and development among different sizes of setts after planting. Similar findings were reported by [Pugalendhi et al. \(2021\)](#). [Pugalendhi & Velmurugan \(2020\)](#) reported that chip bud consists of one to two nodal cuttings planted in protrays recorded the minimum number of days required for sprouting, maximum sprouting percentage, highest length and number of roots

in Cassava. In the present study, tissue cultured plants and six budded setts recorded more vigour in growth with respect of plant height, stem girth and number of leaves. Yield of tubers in cassava is highly variable and is influenced by many factors. The present study indicated that the improvement in plant height, stem girth, number of leaves and length and girth of stem caused increase in tuber yield. This is attributed to the dense foliage which provided the maximum photosynthetic area for the synthesis of assimilates. Similar results were also obtained by Beck (1960). The tuber yield depends largely on the increase of tuber girth which act as a physiological may sink as observed in sweet potatoes by Lowe & Wilson (1975). A positive effect was observed between yield and length and girth of tuber in the present study and is line with that of studies conducted by Magoon et al. (1972) Muthukrishnan et al. (1973) and Pugalendhi et al. (2021). The tuber length/ girth ratio was higher in tissue culture plants and longer setts. Such an inter relationship of girth, weight and length of tubers suggests that the tuber length and girth are the most important traits that could be used as reliable selection indices for high yield as evidenced from this study. In the present study, the plants raised from tissue cultured plantlets recorded significantly higher tuber yield. This is because that the tissue culture plants are observed to be established more quickly, grow more vigorously and produce higher yields. A similar observation was also recorded by Manivasagam et al. (2006). The number of tuberous roots was also more in tissue culture plants while it was less in other treatments because of lower production of fibrous roots. The tissue culture plants recorded the highest yield followed by longer setts. The leaves and petioles had a positive association with the yield of tubers in the normal longer setts also. A similar view was expressed by Estevao et al. (1972). These findings have revealed that the longer the setts also used for planting, the greater the yield expected from them. This was confirmed by Loria (1962), Krochmal (1969), Silva (1971), Mohankumar & Mandal (1971) and Pugalendhi et al. (2021). Since tapioca tubers are used as raw material for many industries, the starch content is an important factor to be reckoned with while assessing the suitability of the tubers for this purpose (Geetha et al., 2021). The increase in starch content was due to the increased production of photosynthates as observed by Sinha & Nair (1968), Geddes et al. (1965), Singh & Mandal (1970) and Resenthal (1972). In the present study, the HCN content was observed in the flesh of the tuber at the harvesting stage. Similar observation was recorded by Moh & Alan (1972) and Muthuswamy et al. (1974). In the present study, among the treatments, all the varieties, the HCN content was below the toxicity level of 50-100mg kg<sup>-1</sup> as postulated by Cock et al. (1979).

## Conclusion

Among the four treatments (different kinds of planting materials) followed in this study, the treatment T<sub>1</sub> Tissue cultured plants recorded a significantly higher number of tuberous roots and ultimately enhanced amount of tuber production. This is mainly due to the quick establishment, vigorous growth and virus free features of tissue culture plantlets. The cassava cv. CO 2 planting materials raised from tissue culture protocols had recorded the lowest incidence of cassava mosaic disease with higher tuber yield.

## Acknowledgment

The authors are highly grateful to the Tamil Nadu Agricultural University for funding and the authors are highly thankful to the Professor and Head, Department of Vegetable Science, TNAU, Coimbatore for technical and logistical support.

## Author contributions

Sundaramoorthy Nanthakumar: Designed and conducted experiments and manuscript writing. Krithika: Conducted experiment, data collection and analysis. Muthusamy Prabhu: Manuscript editing. Govindasamy Ashok Kumar: Statistical analysis.

## Funding

No financial support.

## Conflict of interest

The author declares no conflict of interest. The manuscript has not been submitted for publication in any other journal.

## Ethics approval

Not applicable.

## References

Beck, B. D. (1960). Cassava trials on Moor plantation. Proceedings of third International Symposium on Tropical Crops, Ibadan.

Cock, J. H., Franklin, D., Sandoval, G. & Juri, P. (1979). The ideal cassava plant for maximum yield. *Crop Science*, 1, 271 - 279.

Estevao, E. M., Begazo, J. C. E. O., Brandao, S. S., & Oliveira, L. D. (1972). Production of roots and shoots of cassava varieties and the relationship between characteristics of the above-ground parts and root production. *Revista Ceres*, 19(105), 311 - 327.

Faria, R. T & Illg, R. D. (1995). Micropropagation of *Zingiber spectabile* Griff. *Scientia Horticulturae*, 62, 135 - 137.

Geddes, R., Greenwood, C. R. & Mackenzie, S. (1965). The biosynthesis of starch granules. III. The properties of the components of starches from the growing potato tuber. *Carbohydrate Research*, 1(1), 71 - 82.

Geetha, R., Sankari, A., Pugalendhi, L., Vennila, P., Swarnapriya, R. & Thangamani, C. (2021). Studies on functional properties of cassava var. Yethapur Tapioca-2 for its suitability as an ideal industrial substitute for the grain starches. *Pharma Innovation*, 10(5), 9 - 12.

Hernández - Carmona, F., Morales-Matos, Y., Lambis - Miranda, H., Pasqualino, J. (2017). Starch extraction potential from plantain peel waste. *Journal of Environmental Chemical Engineering*, 5(5), 4980 - 4985.

Indira, P. & Sinha, S. K. (1969). Colorimetric method for determination of HCN in tubers and leaves of Cassava. *Indian Journal of Agricultural Science*, 39, 1021-1023.

Krochmal, A. (1969). Propagation of cassava. *World Crops*, 21(3), 193 - 195.

Loria, M. W. (1962). Influence of size and positioning of cassava cuttings on rooting, yield and foliage production. *Proceedings of Caribbean Region, American Society of Horticultural Science*, 6, 20 - 23.

Lowe, S. B. & Wilson, L. A. (1975). Yield and yield components of six sweet potato (*Ipomoea batatas*) cultivars. Contribution of yield components to tuber yield. *Experimental Agriculture*, 11 (1), 39 - 48.

Lowry, O. H., Rosebrough, N. J., Farr, A. L., & Randall, R. J. (1951). Protein measurement with the Folin phenol reagent. *J biol Chem*, 193(1), 265-275.

Magoon, M. L., Krishnan, R. & Lakshmi, K. (1972). Association of plant and tuber characters with yield of cassava. *Tropical root and tuber crops Newsletter*, 5, 29 - 30.

Manivasagam, S., Rabindran, R., Balasubramanian, P., & Natarajan, S. (2006). Studies on cassava mosaic disease with special reference to detection and virus variability. In *Papers of the 14th Triennial Symposium of International Society for Tropical Root Crops, Central Tuber Crops Research Institute, Sreekariyam, Thiruvananthapuram, Kerala* (p. 153).

Moh, C. C. & Alan, J. J. (1972). The use of the Guignard test for screening cassava cultivars of low HCN content. *Tropical Root and Tuber crops Newsletter*, 6, 29 - 31.

Mohankumar, C. R. & Mandal, R. C. (1971). Studies on sett size and depth of planting of tapioca (H 165). Annual Report, Central Tuber Crops Research Institute. Thiruvananthapuram, Kerala.

Muthukrishnan, C. R., Thamburaj, S., Shanmugam, A. & Subramaniyan, A. S. (1973). Relationship of certain yield components in tapioca (*Manihot esculenta* Crantz). *Madras Agricultural Journal*, 60, 9 - 12.

Muthuswamy, P., Krishnamoorthy, K. K., Muthukrishnan, C. R., Thamburaj, S. & Shanmugam, A. (1974). HCN content of cassava peel as affected by fertilizer application. *Current Science*, 43(10), 312.

Panse, V. G. & Sukhatme, P. V. (1954). Statistical Methods for Agricultural Workers, Second Edition, ICAR, New Delhi.

Pugalendhi, L. & Velmurugan, M. (2020). Standardization of Rapid Multiplication Technique (RMT) in Cassava (*Manihot esculenta* Crantz.). *Int. J. Curr. Microbiol. App. Sci.*, 9(5), 3021- 3025.

Pugalendhi, L., Velmurugan, M., Kavitha, P. S., Kalarani, M. K., Senthil, N., Deivamani, M. & Venkatachalam, S. R. (2021). Evaluation and characterization of high yielding cassava mosaic resistant variety YTP 2. *International Journal of Plant & Soil Science*, 33(22), 198-208.

Radhakrishnan, A. R. S., Suja, G., & Sreekumar, J. (2022). How sustainable is organic management in cassava? Evidences from yield, soil quality, energetics and economics in the humid tropics of South India. *Scientia Horticulturae*, 293, 1-12.

Resenthal. (1972). Characteristics of granules. *Awais da Academica Brasileira de Ciencias.*, 44(1), 55 - 60.

Silva, J. R. D. A. (1971). Cassava research at Agronomic Institute of Sao Paulo state. *Agronomia.*, 23, 49 - 71.

Singh, K. D. & Mandal, R. C. (1970). Studies on the cultural practices in different tuber crops. Plant population in different types of cassava. Annual report, Central Tuber Crops Research Institute, Thiruvananthapuram, Kerala, India, 52 - 54.

Sinha, S. K. & Nair, T. V. R. (1968). Studies on the variability of cyanogenic glucoside content in cassava tubers. *Indian Journal of Agricultural Science*, 38(6), 958 - 963.

Szyniszewska, A. M. (2020). CassavaMap, a fine-resolution disaggregation of cassava production and harvested area in Africa in 2014. *Scientific Data*, 7(159), 1-8.

Tonukari, N. J. (2004). Cassava and the future of starch. *Electronic Journal of Biotechnology*, 7(1), 5-8.