

# Genetic variability and association analysis for yield and yield related traits in finger millet (*Eleusine coracana* (L.) Gaertn)

Ashenafi Yayeh<sup>1\*</sup>, Wossen Tarekegne<sup>2</sup>, Mesfin Abate<sup>1</sup>

<sup>1</sup>Debre Markos University, Department of Plant Sciences, P.O. Box 269, Debre Markos, Ethiopia.

<sup>2</sup>Bahir Dar University, School of plant Sciences, P.O. Box 79, Bahir Dar, Ethiopia.

Received: 01 February 2021

Accepted: 29 May 2021

Published: 30 June 2021

\*Correspondence

Ashenafi Yayeh  
ashe.yayeh@gmail.com

Finger millet (*Eleusine coracana* (L.) Gaertn) is regarded as cereal of hope because of its role in subsistence agriculture, and source of food for millions of poor people in Africa alone. Knowledge of the nature and magnitude of variation existing in breeding materials, interrelationships between quantitatively inherited plant traits is great importance for effective breeding. Forty nine finger millet genotypes were tested at single location at Womberma, West Gojam Zone with the objectives of estimating the genetic variability, association among characters, and to estimate genetic divergence among the genotypes and clustering them in divergent groups. The experiment was conducted using simple lattice design with two replications. Results showed that genotypes had high values of genotypic and phenotypic coefficient of variation for effective tiller per plant, fingers per head, grain yield, biomass yield, lodging and blast severity. High heritability estimated was obtained for all of the traits ranged from 71.43 to 99.56%. Cluster analysis revealed that the 49 genotypes were grouped into nine clusters. Maximum and minimum intra and inter cluster distances was 6.12-16.33 and 17.08-226.28 respectively. Principal component analysis indicated that three principal components explained about 68.07% of the total variation. Differentiation of the genotypes into different cluster was because of accumulative effect of a number of characters rather than small contribution of each individual character.

*Key words:* phenotypic variability, genotypic variability, genetic advance

## INTRODUCTION

Finger millet (*Eleusine coracana* (L.) Gaertn) is a tetraploid ( $2n=4x=36$ ) is highly self-pollinating crop belongs to the family Poaceae, and it is a small seeded cereal crop grown in moisture stressed areas of the semi-arid tropics of the world, especially in East Africa, India and in other Asian countries (Rudin et al., 2004). Ethiopia is the centre of diversity for finger millet. It is mainly grown in northern, north western and south western part of the country. Its annual world production was about 30.5 million tons: 12.4 million tons were produced in Africa mainly eastern and southern African countries (FAOSTAT, 2015). Finger millet is produced in different regional states of Ethiopia: such as Benshangul-Gumuz, Tigray, SNNP, Amhara and Oromia Regions. Total area coverage of finger millet in Ethiopia was 446,909.00 ha with a total production and productivity of 1.035 million tons 2.317 tons per ha, respectively (CSA, 2019). It has large area coverage in the Amhara region as compared to others, with 248,292.40 ha of land and giving 0.571 million tons production (CSA,

2019). Finger millet is highly nutritious and its grain contains 0.3-0.4% calcium, 2.5-3.5% minerals, 5-8% protein, 15-20% dietary fibre and 65-75% carbohydrates (Chetan and Malleshi, 2007). The availability of diverse genetic resources is a prerequisite for genetic improvement of any crop including finger millet. Besides the availability of genetic resources, their characterization is essential for effective utilization in crop improvement programs. Success of hybridization programme depends to a large extent upon the choice of suitable parents of diverse origin with the possibility of obtaining large frequency of transgressive segregants (Kumar et al., 2010). Meanwhile the objective of any crop breeding programs, selection is one of an integral part which genotypes with high productive in a given environment could be developed. Selection for high yield is made difficult by the complex nature of this trait. Yield per unit area is the end product of components of several yield contributing characters (Singh and Singh, 1973). The polygenic

inheritance of yield related traits makes selection more difficult. Moreover, these complex traits are highly influenced by environment, which reduces the progress to be achieved through direct selection. In such cases, there is another option to hasten the genetic improvement which is known as indirect selection for yield. Yield related traits show association among themselves and with yield. Plant breeder have to find significant correlations among yield and yield related traits, and effect of yield component traits on grain yield to predict the superior cross combinations and to select ideal plant type with increased yield (Keerthana et al., 2019). In Ethiopia there is limited information on the extent and pattern of variability of finger millet a collection under diversified agro-climatic conditions (Kassahun and Solomon, 2017). So the objective of the study (1) to estimate the magnitude of genetic diversity for yield; yield related traits of finger millet genotypes (2) to estimate the genotypic and phenotypic association among characters (3) to determine the direct and indirect effect of yield related traits on seed yield, and (4) to clustering of genotype based on genetic divergence.

## MATERIALS AND METHODS

### Description of the Study Site

The field experiment was conducted during the 2019/20 main cropping season in the Womberma District, North West Ethiopia. Womberma is located at longitude 36 56'32"E and latitude 10 38' 37"N in northern highlands of Gojam in Ethiopia with maximum and minimum altitude of 2125 and 783 meters above sea level (m.a.s.l.). The annual rainfall and temperatures of the area maximum and minimum 1430 mm, 1100 mm and 24°C, 14 °C, respectively.

### Experimental materials

The experimental materials consisted of 48 finger millet genotypes and one local varies as a check obtained from Ethiopian Institute of Biodiversity Institute (EBI) and in the study area, respectively,

**Table 1. List of finger millet genotypes used in the study arranged according to their collection regions and zones**

| NO. | Genotypes  | Source | regional state | Place of Collection |                  |                     |
|-----|------------|--------|----------------|---------------------|------------------|---------------------|
|     |            |        |                | Zone                | Woreda/ District | Altitude (m.a.s.l.) |
| 1   | ACC#215829 | EBI    | Amhara         | West Gojam          | Jabi tehnan      | 1990                |
| 2   | ACC#215837 | EBI    | Amhara         | West Gojam          | Jabi tehnan      | 2050                |
| 3   | ACC#215847 | EBI    | Amhara         | West Gojam          | Bahir dar        | 1820                |
| 4   | ACC#215848 | EBI    | Amhara         | West Gojam          | Bahir dar        | 1820                |
| 5   | ACC#215876 | EBI    | Amhara         | West Gojam          | Adet             | 2250                |
| 6   | ACC#215878 | EBI    | Amhara         | West Gojam          | Adet             | 2320                |
| 7   | ACC#215887 | EBI    | Amhara         | East gojam          | Hu.j enese       | 1880                |
| 8   | ACC#215890 | EBI    | Amhara         | West Gojam          | B.wemberma       | 2080                |
| 9   | ACC#215911 | EBI    | Amhara         | Agew awi            | Dangela          | 2200                |
| 10  | ACC#215913 | EBI    | Amhara         | Agew awi            | Dangela          | 2110                |
| 11  | ACC#215967 | EBI    | Amhara         | South gondar        | Fogera           | 1880                |
| 12  | ACC#215968 | EBI    | Amhara         | North gondar        | Gond.zuria       | 2500                |
| 13  | ACC#215973 | EBI    | Amhara         | North gondar        | Gond. zuria      | 2090                |
| 14  | ACC#215975 | EBI    | Amhara         | North gondar        | Dembia           | 1980                |
| 15  | ACC#215976 | EBI    | Amhara         | North gondar        | Dembia           | 1860                |
| 16  | ACC#215984 | EBI    | Amhara         | North gondar        | Alefa            | 1870                |
| 17  | ACC#215995 | EBI    | Amhara         | South gondar        | Farta            | 2330                |
| 18  | ACC#215996 | EBI    | Amhara         | South gondar        | Farta            | 2330                |
| 19  | ACC#225891 | EBI    | Amhara         | South gondar        | Este             | 2360                |
| 20  | ACC#225894 | EBI    | Amhara         | West Gojam          | Merawi           | 1900                |
| 21  | ACC#225896 | EBI    | Amhara         | West Gojam          | Merawi           | 1920                |
| 22  | ACC#228307 | EBI    | Amhara         | West Gojam          | B.wemberma       | NA                  |
| 23  | ACC#235834 | EBI    | Amhara         | North gondar        | La. armacho      | 1920                |
| 24  | ACC#237443 | EBI    | Amhara         | North wello         | Guba lafto       | 2100                |
| 25  | ACC#238338 | EBI    | Amhara         | North gondar        | Teda             | 1900                |
| 26  | ACC#242106 | EBI    | Amhara         | West Gojam          | Achefer          | 1955                |
| 27  | ACC#242107 | EBI    | Amhara         | West Gojam          | Achefer          | 2020                |
| 28  | ACC#242135 | EBI    | Amhara         | North wello         | Guba lafto       | 1910                |

Table 1. Continue...

| NO. | Genotypes   | Source | regional state | Place of Collection  |                  |                     |
|-----|-------------|--------|----------------|----------------------|------------------|---------------------|
|     |             |        |                | Zone                 | Woreda/ District | Altitude (m.a.s.l.) |
| 29  | ACC#243616  | EBI    | Amhara         | South wello          | Kalu             | 2290                |
| 30  | ACC#243617  | EBI    | Amhara         | South wello          | Kalu             | 1780                |
| 31  | ACC#216046  | EBI    | Oromiya        | West wellega         | Nejo             | 1680                |
| 32  | ACC#216048  | EBI    | Oromiya        | West wellega         | Mana sibu        | 1640                |
| 33  | ACC#216049  | EBI    | Oromiya        | West wellega         | Mana sibu        | 1600                |
| 34  | ACC#216051  | EBI    | Oromiya        | West wellega         | Lalo asabi       | 1910                |
| 35  | ACC#216054  | EBI    | Oromiya        | West wellega         | Ayra guliso      | 1730                |
|     | ACC#24392   | EBI    | Oromiya        | West wellega         | Gimbi            | 1780                |
| 37  | ACC#212462  | EBI    | Oromiya        | West harerge         | Habro            | 1400                |
| 38  | ACC#24393   | EBI    | Oromiya        | West harerge         | Lalo asabi       | 1745                |
| 39  | ACC#24394   | EBI    | Oromiya        | West harerge         | Lalo asabi       | 1730                |
| 40  | ACC#215804  | EBI    | Oromiya        | West wellega         | Sayo             | 1950                |
| 41  | ACC#27885   | EBI    | SNNP           | Bench maji           | Debubbench       | 1400                |
| 42  | ACC#227973  | EBI    | SNNP           | Hadiya               | Badawacho        | NA                  |
| 43  | ACC#227974  | EBI    | SNNP           | Kembata alabana temb | Alaba            | NA                  |
| 44  | ACC#227975  | EBI    | SNNP           | Hadiya               | Limo             | NA                  |
| 45  | ACC#240506  | EBI    | SNNP           | Kembata alabana temb | Alaba            | NA                  |
| 46  | ACC#241769  | EBI    | SNNP           | Keficho shekicho     | Chena            | 1500                |
| 47  | ACC#244798  | EBI    | SNNP           | Bench maji           | Konso special    | 2169                |
| 48  | ACC#229722  | EBI    | BG             | Metekel              | Dangur           | 1750                |
| 49  | Local check |        |                |                      |                  |                     |

NA= Not Available, EBI = Ethiopia Biodiversity Institute, SNNP= South Nation and Nationality of People, BG= Benishangul Gumuz, M.A.S.L. =Meter above sea level.

#### Experimental design and trial management

The trial was conducted using 7 X 7 simple lattice designs with two replications Gomez and Gomez (1984). Each experimental plot has an area of 1.6 m<sup>2</sup> with two rows of 2 m length space of 0.40 m between rows, 0.15 m between plants, distance between plot and block was 0.5m and 1m respectively. NPSB and urea were applied at the rate of 100 and 50 kg/ha, respectively, NPSB was applied at planting and urea was applied in split; half at planting and the rest half at the time of tillering. Thinning was done after three weeks of planting to maintain the space between plants and to balance the plant density. Other crop management practices were applied following the recommended practices.

#### Data Collection and Sampling procedures

Data were recorded on 13 quantitative traits on plot and plant basis using descriptors for finger millet (IBPGR, 1985). Ten representative plants were randomly selected from the middle rows of each plot. Quantitative data such as plant height, number of effective tillers per plant, number of ears per plant, number of fingers per ear and finger length were collected on per plant basis using ten randomly selected plants in each plot. data such as ,days to flowering, days to maturity, biomass yield per plot, grain

yield per , harvest index, thousand kernel , lodging index and blast diseases were collected on the whole plot basis.

#### Data Analysis

Analysis of variance was done using the procedures outlined by Gomez and Gomez (1984) with the help of SAS, version 9.2 (SAS Institute Inc.). Duncan's Multiple Range Test (DMRT) was used for mean separation at 1% and 5% probability level. Cluster analysis based on Tocher's method and correlation matrix based principal component analysis was conducted using SAS- JMP Software. The means were used for cluster and PCA analysis and calculating the genetic distance between groups. The phenotypic ( $\sigma^2_p$ ), genotypic ( $\sigma^2_g$ ) and error ( $\sigma^2_e$ ) variances were calculated from expected mean squares of analysis of variance and genetic advance and genetic advance as percent of mean (GAM) were also estimated.

1. Genotypic variance ( $\delta^2_g$ ) =  $\frac{MSG - MSe}{r}$  (Burton and De Vane, 1953)
2. Phenotypic variance ( $\delta^2_p$ ) =  $\delta^2_g + \delta^2_e$  (Burton and De Vane, 1953)

3. Genotypic coefficient of variation (GCV) =  $\frac{\sqrt{\hat{\sigma}_g^2}}{x} \times 100$  (Singh and Chaudhury, 1985)
4. Phenotypic coefficient of variation (PCV) =  $\frac{\sqrt{\hat{\sigma}_p^2}}{x} \times 100$  (Singh and Chaudhury, 1985)
5. Broad sense heritability ( $h^2$ ) =  $\frac{\hat{\sigma}_g^2}{\hat{\sigma}_p^2} \times 100$  (Falconer and Mackay (1996)
6. Genetic advance (GA) =  $GA = K\sigma_P h^2$  (Johnson *et al.* 1955 and Allard 1960)
7. Genetic advance as percent of mean (GAM) =  $\frac{GA}{x} \times 100$  (Johnson 1955 and Allard 1960)

## RESULTS AND DISCUSSION

### Analysis of Variance

The Analysis of Variance (ANOVA) revealed highly significant differences ( $P < 0.01$ ) among the accessions for all 13 quantitative (Table 1), indicating that the presence of high genetic diversity among finger millet

953.1 kg per ha (ACC#237443) to 5234.4 kg per ha (ACC#235834) with a mean of 2895.18 kg per ha and that of the check was 2343.8 kg per ha. Number of fingers per ear ranged from 5.25 (ACC#215848) to 14.15 (ACC#244798) with a mean of 8.01 fingers per ear and the check was 6.5. Finger length of the test varieties varied from 6.08 cm (ACC#212462) to 15.95cm (ACC#215973) with mean of 10.69cm while that of the check finger was 9.715cm. Number of effective tillers per plant ranged from 5.2 (ACC#227975) to 14.80 (ACC#215804) with a mean of 9.22 while the check had 7.45 (Table 2).

### Estimates of variance components, heritability and genetic advance

#### Genotypic and phenotypic variance

Estimates of genotypic variance ( $\sigma^2_g$ ), phenotypic variance ( $\sigma^2_p$ ) and error variance ( $\sigma^2_e$ ) variability are presented on (Table 2). The levels of diversity among the accessions were estimated based on the genetic and phenotypic coefficient of variation. Phenotypic coefficient of variability (PCV) values ranged from 4.33 % days to maturity to 97.65% for finger blast diseases, whereas the genotypic coefficient of variability

**Table 2. Mean square values and coefficient of variations in agronomic traits of finger millet**

| Traits         | Mean Squares            |                  |               |       |                  |
|----------------|-------------------------|------------------|---------------|-------|------------------|
|                | Replication block(df=1) | Genotype (df=48) | Error (df=36) | CV %  | R <sup>2</sup> % |
| DH             | 2                       | 97.27**          | 0.78          | 0.91  | 99.4             |
| DM             | 0.65                    | 37.43**          | 5.15          | 1.6   | 91.9             |
| BMV(kg per ha) | 8418733                 | 24773926**       | 561152        | 5.46  | 98.6             |
| GY (kg per ha) | 907026.51               | 1035749.54**     | 62672.11      | 8.65  | 96.2             |
| PH (cm)        | 0.30                    | 307.23676**      | 9.14          | 3.8   | 98.07            |
| FL (cm)        | 0.24                    | 3.2060222**      | 0.12          | 3.23  | 97.7             |
| NEPP           | 1.62                    | 8.583839**       | 0.68          | 8     | 95.82            |
| NFPH           | 1.3                     | 6.8359651**      | 0.15          | 4.84  | 98.87            |
| NETPP          | 0.297                   | 8.4030867**      | 0.15          | 4.21  | 98.86            |
| HI             | 3.42                    | 40.36**          | 2.41          | 7.17  | 96.38            |
| TSW            | 0.03                    | 0.12**           | 0.03          | 6.29  | 86.49            |
| BS             | 48.58                   | 664.84**         | 13.43         | 18.95 | 98.6             |
| LO             | 1.23                    | 774.48**         | 1.67          | 3.36  | 99.85            |

DF= Degree of freedom, CV=Coefficient of variation, R<sup>2</sup>= coefficient of determination, PH= Plant height, NETPP=Number of effective tillers per plant, NEPP=Number of Ears per plant, FL=Finger length, DH=Days to 50% Heading, DM=Days to 50% maturity, BMV=Biomass Yield per hectare, GY= Grain Yield per hectare, HI= Harvest index, TSW= Thousand seed weight, LO=Lodging susceptibility %, BS= blast diseases, \*=Significant at probability level of 0.05 and \*\*=Significant at probability level of 0.01.

accessions which can be exploited through selection. Such differences among finger millet accessions were previously reported by Damot *et al.* (2019 and Yaregal *et al.* (2019).

### Estimates of Mean and Range

The 49 finger millet genotypes showed wide range of variability for all of the characters (Table 2). Days to heading ranged from 87 for (ACC#237443) to 114 days for (ACC#216049) with a mean of 97.9, whereas the local check had 94 days. Days to maturity ranged from 134.5 for (ACC#243616) to 152.5 (ACC #216051) with a mean of 141.35 compared to the check 136.5 days. Grain yielding ability ranged from

(GCV) ranged from 4.02 % days to maturity to 90.76% finger blast diseases. The phenotypic variance was greater than the genotypic variance for all quantitative traits in this study (Table 2). The highest GCV values were obtained from harvest index (20.15%), effective tiller per plant (22.04%), finger per head (22.8%), grain yield (24.98%), biomass yield (25.37%), lodging (51.08%) and finger blast diseases (90.76%), Moderate GCV values were obtained from number of ears per plant (19.28%), Plant height (15.35%), and finger length (11.61% and Low GCV value was recorded for days of maturity (4.02%), Days of heading (7.09%), and thousand seed weigh (8.47%).The high GCV values of these characters suggest that the possibility of improving these trait through direct selection.

**Table 3. Estimates of range, mean, genetic components of variance, heritability and genetic advance**

| No. | Traits | Range  |          | Mean     | SEM(±) | $\sigma^2_p$ | Variance     |              |       | Coefficient of variation |           |         |        |  |
|-----|--------|--------|----------|----------|--------|--------------|--------------|--------------|-------|--------------------------|-----------|---------|--------|--|
|     |        | Min    | Max      |          |        |              | $\sigma^2_g$ | $\sigma^2_e$ | PCV%  | GCV%                     | $h^2$ (%) | GA      | GAM    |  |
| 1   | DH     | 87.00  | 114.00   | 97.90    | 0.88   | 49.025       | 48.245       | 0.78         | 7.15  | 7.09                     | 98.40     | 14.21   | 14.52  |  |
| 2   | DM     | 134.50 | 152.00   | 141.35   | 2.26   | 37.45        | 32.30        | 5.13         | 4.33  | 4.02                     | 86.25     | 10.89   | 7.70   |  |
| 3   | BMY    | 5410.2 | 23628.90 | 13712.40 | 749.1  | 12667539     | 12106387     | 561152       | 25.96 | 25.37                    | 95.57     | 7017.25 | 51.17  |  |
| 4   | GY     | 953.1  | 5234.40  | 2895.18  | 250.34 | 549211       | 486539       | 62672.1      | 25.60 | 24.09                    | 88.59     | 1354.42 | 46.78  |  |
| 5   | PH     | 54.07  | 113.38   | 79.53    | 3.02   | 158.19       | 149.05       | 9.14         | 15.81 | 15.35                    | 94.22     | 24.45   | 30.74  |  |
| 6   | FL     | 6.08   | 14.95    | 10.69    | 0.35   | 1.66         | 1.54         | 0.12         | 12.05 | 11.61                    | 96.34     | 2.56    | 23.95  |  |
| 7   | NEPP   | 5.80   | 16.25    | 10.31    | 0.83   | 4.63         | 3.95         | 0.68         | 20.87 | 19.28                    | 85.31     | 3.79    | 36.72  |  |
| 8   | FPH    | 5.25   | 14.15    | 8.01     | 0.39   | 3.50         | 3.35         | 0.15         | 23.38 | 22.80                    | 95.51     | 3.68    | 45.99  |  |
| 9   | NETP   | 5.20   | 14.80    | 9.22     | 0.39   | 4.28         | 4.13         | 0.15         | 22.43 | 22.04                    | 96.50     | 4.12    | 44.67  |  |
| 10  | HI     | 12.70  | 34.16    | 21.62    | 1.55   | 21.38        | 18.98        | 2.40         | 21.39 | 20.15                    | 88.77     | 8.47    | 39.16  |  |
| 11  | TSW    | 1.93   | 3.20     | 2.64     | 0.17   | 0.07         | 0.05         | 0.03         | 10.02 | 8.47                     | 71.43     | 0.39    | 14.76  |  |
| 12  | BS     | 5.00   | 82.50    | 19.34    | 3.66   | 356.71       | 308.13       | 48.58        | 97.65 | 90.76                    | 92.93     | 35.73   | 184.75 |  |
| 13  | Lo     | 19.00  | 83.50    | 38.48    | 1.29   | 388.08       | 386.41       | 1.67         | 51.19 | 51.08                    | 99.56     | 40.46   | 105.15 |  |

DF= Days of heading, DM=Days to 50% maturity, BMY=Biomass yield, PH= Plant height, FL=Finger length, NEPP=Number of Ears per plant, NFPH= Number of finger per head, NETPP=Number of effective tillers per plant, HI= Harvest index, TSW= Thousand seed weight, BS= blast diseases, LO=Lodging susceptibility%, GY=Grain yield, SE±=Standard error,  $\sigma^2_p$  =Phenotypic variance,  $\sigma^2_g$  =Genotypic variance,  $\sigma^2_e$ =Error Variance, PCV=Phenotypic coefficient of variation, GCV=Genotypic coefficient of variation,  $h^2$  =Broad sense heritability, GA=Expected genetic advance and GAM= Genetic advance as percent of the mean.

#### Estimation of heritability in broad sense and genetic advance

High GCV values were also obtained by Keerthana et al. (2019) for grain yield (42.29%), biomass yield (30.14%), and effective tiller per plant (27.89%). Yaregal et al. (2019) also reported high GCV value for grain yield (26.8%), blast diseases (30.23%) and lodging (68.03%). Moderate GCV values were obtained from number of ears per plant (19.28%), Plant height (15.35%), and finger length (11.61%). In line with Yaregal et al. (2019) for plant height (11.84), Keerthana et al. (2019) and Kassahun and Solomon (2017) for plant height and finger length, but ear per plant was in contrast with reported by (Kassahun and Solomon, 2017). Low GCV value was recorded for days of maturity (4.02%), Days of flowering (7.09%), and thousand seed weigh (8.47%). Similar results were also obtained by Kassahun and Solomon (2017) for days of flowering and days to maturity, Keerthana et al. (2019, Yaregal et al. (2019) and Singamsetti et al. (2018) for days of maturity, Damot et al. (2019) for days of flowering, Devaliya et al. (2018) for thousand seed weight.

Phenotypic coefficient of variability (PCV) values ranged from 4.33 % days to maturity to 97.65% for finger blast. High phenotypic coefficient of variations (PCV) were recorded for Grain yield (25.6%), biomass yield (25.96%), and finger per head (23.38%), effective tiller per plant (22.43%), harvest index (21.39%), finger blast diseases (97.65%) and lodging (51.19%). Similar results were also obtained by, Yaregal et al. (2019) for grain yield, effective tiller per plant, blast diseases and lodging. Keerthana et al. (2019) for grain yield, biomass yield and effective tiller per plant. Number of ears per plant (20.89%), Plant height (15.81%), and finger length (12.05%), thousand seed weight (10.2%) showed relatively moderate PCV. Similar results were also obtained by Keerthana et al. (2019), Yaregal et al. (2019) and Kassahun and Solomon, (2017) for plant height and finger length; but ear per plant was in contrast with the report of Kassahun and Solomon (2017). Low PCV values were recorded for days to flowering (7.15%), days to maturity (4.33%). In Similar studies were reported by Kassahun and Solomon (2017) for days of flowering and days to maturity, Keerthana et al. (2019) and Singamsetti et al. (2018) for days to maturity and by Damot et al. (2019) for days of flowering.


Estimates of heritability in broad sense ranged from 71.43% for thousand seed weight to 99.56% for lodging (Table 2). Heritability estimate was high (>80%) days to flowering (98.4%), days to maturity (86.25%), biomass yield (95.57%), grain yield (88.59%), plant height (94.22%), finger length (96.34%), ear per plant (85.31%), finger per head (95.51%), effective tiller per plant (96.5%), harvest index (88.77%), finger blast diseases (92.93%) lodging susceptibility (99.56%), indicating that the variation observed were mainly under genetic control and were less influenced by the environment and the possibility of progress from selection. Moderate heritability was recorded for thousand seed weight (71.43%). This result is in line with the finding of Damot et al. (2019) for days to maturity (72.45%), grain yield (76.47%), biomass yield (83.17%), ear per plant (92.03%) and plant height (93.35%); Yaregal et al. (2019) for finger length (66.12), days of heading (95.46), and lodging (86.41); Keerthana et al. (2019) days to flowering (97.98), days to maturity (98.14), biomass yield (97.29), grain yield (96.49), plant height (98.95), finger length (74.46), finger per head (97.37), thousand seed weight (89.56, but finger blast diseases is contrast to the finding of (Keerthana et al., 2019). The highest expected genetic advance as percent of mean from selection of the top 5% of the accessions was obtained for grain yield was 1354.42 kg per ha indicating that whenever we select the best 5% high yielding genotypes as parents, mean grain yield of progenies could be improved by 1354.42 kg per ha, that is, mean genotypic value of the new population for grain yield were improved from 2895.18 kg per ha to 4249.6 kg per ha. In the same way, it was 14.1 for number of ears per plant, 20729.65 kg per ha for biomass yield, 30.09% for harvest index, 13.25cm for finger length, 13.34 for effective tiller and 11.69 for number of fingers per ear (Table 2). Maximum genetic advance as Table 4. Eigenvectors and eigenvalues of the first three principal components (PCs) for 13 characters of finger millet genotypes percent of mean (GAM) at 5% selection intensity was recorded for finger blast severity (184.75%) and lodging susceptibility (105.1%) followed by biomass yield (51.17%). It was low for days of maturity (7.70%) and days of heading (14.52%) (Table 3).

**Table 4. Eigenvectors and eigenvalues of the first three principal components (PCs) for 13 characters of finger millet genotypes**

|                           | PCA1         | PCA2         | PCA3         |
|---------------------------|--------------|--------------|--------------|
| <b>Eigen Value (Root)</b> | <b>4.651</b> | <b>2.427</b> | <b>1.772</b> |
| <b>% Var. Exp.</b>        | <b>35.77</b> | <b>18.67</b> | <b>13.63</b> |
| <b>Cum. Var. Exp.</b>     | <b>35.77</b> | <b>54.4</b>  | <b>68.07</b> |
| DH                        | 0.31696      | -0.27122     | -0.14021     |
| DM                        | 0.29029      | 0.0344       | 0.26354      |
| BMY(kg per ha)            | 0.39191      | -0.06981     | 0.19323      |
| GY(kg per ha)             | 0.37868      | 0.20425      | -0.20084     |
| PH(cm)                    | 0.34973      | -0.18102     | 0.30367      |
| FL(cm)                    | 0.29414      | 0.13128      | 0.11061      |
| NEPP                      | -0.06631     | 0.49291      | 0.3099       |
| NFPH                      | 0.1748       | -0.46531     | -0.05577     |
| NETPP                     | 0.24191      | 0.40641      | 0.14007      |
| HI                        | -0.01859     | 0.3493       | -0.52355     |
| TSW                       | 0.27715      | 0.22989      | -0.29813     |
| BS                        | -0.372       | 0.034        | 0.24792      |
| LO                        | -0.02076     | 0.16734      | 0.43075      |

DH= Days of heading, DM=Days to 50% maturity, BMY=Biomass yield, GY= Grain yield, PH= Plant height, FL=Finger length, NEPP=Number of Ears per plant, NFPH=Number of finger per head, NETPP=Number of effective tillers per plant, HI= Harvest index, TSW= Thousand seed weight and BS= blast diseases, LO= Lodging index.

**Figure 1. Dendrogram of 49 genotypes of finger millet based on evaluation for 13 characters**



The traits with high expected genetic advance values were biomass yield (51.17%), grain yield (46.78%), finger per head (45.99%), effective tiller (44.67%), ear per plant (36.72%), plant height (30.74%). Moderate GAM estimates were obtained for finger length (23.95%), thousand seed weight (14.76%), days of flowering (14.52%); and day of maturity (7.70%) showed comparatively low values of genetic advance expressed as percent of the mean (Table 3). Similar results reported by Devaliya et al. (2018) for biomass yield, grain yield, effective tiller per plant had high heritability and genetic advance as per cent of mean, and Moderate genetic advance as per cent of mean was recorded for number of fingers per ear, thousand seed weight and days to 50% flowering, while the remaining characters had shown low genetic advance as expressed as percentage of mean.

#### Principal component analysis

The principal component analysis (PCA) was done to identify the critical trait which abetted selection for designing future breeding strategies and

recognizes which trait explained more of variation out of 13 traits of finger millet genotypes. For this finding, the three principal components possessed eigenvalues greater than one considered important for explaining the variations observed in the genotypes. These principal component analyses revealed that three principal components PC1 to PC3 with eigenvalues, 4.651, 2.427 and 1.772 respectively, have accounted for 68.07% of the total variation (Table 4). The three principal components PC1, PC2 and PC3 with values of 35.77%, 18.67% and 13.63%, respectively, contributed more to the total variability Characters having relatively higher value in the first principal component contributed maximum towards variability (35.77). Like plant height, days of flowering, days of maturity, fingers length, biomass yield and grain yield, blast diseases had more contribution to the total diversity and they were the ones that most differentiated the clusters. The second principal component (PC2) described percent of total variance and the characters viz., number of productive tillers, No. of fingers per head, Harvest index and Thousand seed weight showed the maximum variance in this

**Table 5. Mean value of 13 quantitative characters for the nine clusters of 49 finger millet genotypes**

| Traits         | cluster  |          |          |        |          |          |          |          |         |
|----------------|----------|----------|----------|--------|----------|----------|----------|----------|---------|
|                | I        | II       | III      | IV     | V        | VI       | VII      | VIII     | IX      |
| DF             | 107.86   | 98.00    | 108.25   | 98.17  | 91.21    | 91.75    | 98.50    | 98.83    | 90.33   |
| DM             | 147.14   | 142.64   | 141.88   | 136.58 | 138.57   | 143.13   | 140.88   | 140.67   | 136.67  |
| BMV(kg per ha) | 16938.10 | 15366.00 | 19190.00 | 10247  | 11830.00 | 14385.00 | 13525.00 | 11167.00 | 7347.70 |
| GY (kg per ha) | 3872.54  | 2825.40  | 3396.10  | 2999.7 | 2813.20  | 2955.10  | 2683.60  | 1875.00  | 1234.40 |
| PH (cm)        | 92.10    | 79.80    | 100.84   | 63.56  | 68.05    | 81.81    | 88.61    | 80.40    | 60.91   |
| FL (cm)        | 11.96    | 11.64    | 10.56    | 9.64   | 10.21    | 10.96    | 11.26    | 10.19    | 7.95    |
| NEPP           | 10.49    | 9.61     | 6.74     | 8.68   | 11.16    | 13.53    | 10.22    | 9.70     | 9.68    |
| NFPH           | 8.28     | 9.09     | 12.56    | 7.14   | 6.19     | 6.74     | 7.20     | 9.77     | 7.60    |
| NETPP          | 10.24    | 8.41     | 10.35    | 8.16   | 9.56     | 11.64    | 8.54     | 6.38     | 5.80    |
| HI             | 23.33    | 18.68    | 17.97    | 29.68  | 23.85    | 20.94    | 19.90    | 16.82    | 17.04   |
| TSW            | 2.88     | 2.77     | 2.63     | 2.68   | 2.60     | 2.69     | 2.44     | 2.21     | 2.29    |
| BS             | 7.43     | 10.27    | 9.13     | 10.25  | 28.85    | 14.75    | 14.25    | 55.83    | 60.50   |
| LO             | 31.29    | 27.71    | 32.88    | 34.00  | 27.14    | 61.38    | 67.75    | 38.33    | 23.33   |

DF=Days of 50% flowering, DM=Days to 50% maturity, BMV=Biomass yield, GY= Grain yield, PH= Plant height, FL=Finger length, NEPP=Number of Ears per plant, NFPH=Number of finger per head, NETPP=Number of effective tillers per plant, HI= Harvest index, TSW= Thousand seed weight and BS= blast diseases, LO= Lodging index.

**Table 6. Average intra (bold) and inter cluster (off diagonal) D<sup>2</sup> values among eight clusters in 49 finger millet genotypes**

| Cluster | I            | II           | III         | IV           | V            | VI           | VII         | VIII        | IX          |
|---------|--------------|--------------|-------------|--------------|--------------|--------------|-------------|-------------|-------------|
| I       | <b>14.29</b> | 31.992**     | 48.45**     | 116.423**    | 86.274**     | 41.929**     | 27.995**    | 67.9**      | 201.815**   |
| II      |              | <b>14.29</b> | 58.153**    | 60.023**     | 29.914**     | 23.503*      | 20.068      | 32.473**    | 97.778**    |
| III     |              |              | <b>8.16</b> | 135.817**    | 127.525**    | 95.526**     | 71.975**    | 71.184**    | 226.284**   |
| IV      |              |              |             | <b>12.24</b> | 34.401**     | 106.204**    | 84.844**    | 88.362**    | 99.336**    |
| V       |              |              |             |              | <b>14.29</b> | 46.496**     | 46.621**    | 40.442**    | 38.984**    |
| VI      |              |              |             |              |              | <b>16.33</b> | 17.057      | 61.599**    | 125.314**   |
| VII     |              |              |             |              |              |              | <b>8.16</b> | 38.836**    | 123.776**   |
| VIII    |              |              |             |              |              |              |             | <b>6.12</b> | 64.741**    |
| IX      |              |              |             |              |              |              |             |             | <b>6.12</b> |

$c^2 = 21.03$ , and  $26.22$  at 5% and 1% probability level respectively

\*, \*\* Indicates significance at 5% and 1% level, respectively.

principal component. The third principal component (PC3) was characterized by 13.63 per cent contribution towards the total variability. The characters viz., plant height, number of ear per plant, harvest index, lodging, contributed maximum variance in this principal component. In line with this study (Yaregal et al., 2019) using 24 germplasm accessions of finger millets reported three clusters, (Kassahun And Solomon, 2017) using 60 germplasm accessions of finger millets reported three clusters and (Patel et al., 2017) found three PCs using 65 finger millet germplasm accessions. In contrast to this study, (Harshal et al., 2017) using 65 germplasm accessions of finger millets reported five clusters and (Damot et al., 2019) using 225 germplasm accessions of finger millets reported five clusters.

#### Cluster analysis

The D<sup>2</sup> values based on the mean of genotypes resulted in classifying the 49 finger millet genotypes into nine distinct clusters is shown in (Table 5). Around 45% of the genotypes were grouped under cluster I, II and V consisting of 7 genotypes each and lowest 12% of the genotypes were grouped under cluster VIII and IX consisting of 3 genotypes each. clusters I (14.29%) was the second largest cluster containing 7 genotypes, cluster II (14.29%) consisting 7, cluster III (8.16%) with 4 genotypes, cluster IV (12.24%) with 6 genotypes, cluster V (14.29%) had 7, Cluster VI (16.33%) was the largest cluster containing 8 genotypes, cluster VII (8.16%) with 4 genotypes, cluster VIII (6.12%) had 3 genotypes and cluster IX (6.12%) with 3 genotypes (Table 6). The

usefulness and of success Mahalanobis' D<sup>2</sup> analysis in genetic divergence in finger millet has been studied by Kandel et al. (2019), Yaregal et al. (2019) and Damot et al. (2019).

#### CONCLUSION

The result of the current study showed that ample genetic diversity existed among finger millet accessions to be used in future breeding program through selection and hybridization. Wide ranges of genetic diversity were observed in most of quantitative traits. The presence of genetic diversity between the accessions and the range of variation showed the chance of genetic improvement via selection and/or crossing. High heritability accompanied with high genetic advance was estimated for number of ears per plant, number of finger per plant, finger length, days to flowering, grain yield, biomass yield, lodging susceptibility and blast severity. Hence, these traits, which showed high heritability values coupled with high genetic advance as percent of means, indicated the possibility to improve finger millet through selection. Therefore, the existence of high genetic diversity is a basis for comprehensive and systematic germplasm collections of finger millet for further genetic conservation and utilization.

#### AUTHOR CONTRIBUTIONS

This work was carried out in collaboration between authors. Author Ashenafi Yaheh conducted experiments, collected data and performed

statistical analysis and drafted the manuscript. Author Wossen performed statistical analysis and drafted the manuscripts. Mesfin Abate has given his inputs for the generation of this manuscript. All authors read and approved the final manuscript.

## ACKNOWLEDGMENTS

First of all, I would like to thank Lord Jesus for being with us in all aspects during this work. I am grateful to my major advisor, Dr. Wossen Tarekegne for his encouragement which I enjoyed and thorough review and constructive criticisms and suggestions on the manuscript that enable me to shape up this thesis in its final form. My special appreciation goes to my co-advisor, Dr. Mesfin Abate for his valuable comments in the course of preparation of the proposal as well as this manuscript. I would like to thank the Womberma woreda agriculture office for facilitating budget I would like to express my appreciation for staff members of Womberma woreda agriculture office especially, facilitating for vehicle for field supervision and data collection. I would like to thank the Ethiopian Biodiversity Institute for the supply of seed of finger millet genotypes.

## COMPETING INTERESTS

The authors declare that they have no competing interests.

## ETHICS APPROVAL

Not applicable.

## REFERENCES

- Allard, R.W., (1960). Principles of plant breeding. John Willey and Sons, New York. 485.
- Anteneh, D., Mekbib, F., Tadesse, T., & Dessalegn, Y. (2019). Genetic Diversity among Lowland Finger Millet (*Eleusine coracana* (L.) Gaertn) Accessions. *Ethiopian Journal of Agricultural Sciences*, 29(2), 93-108.
- Devaliya, S.D., Manju Singh, C. G. Intawala & Ranjit N. Bhagora, (2018). Genetic Variability Studies in Finger Millet (*Eleusine coracana* (L.) Gaertn). *Int. J. Pure App. Biosci.*, 6 (1): 1007-1011.
- Burton, G. W., & Devane, D. E. (1953). Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material 1. *Agronomy Journal*, 45(10), 478-481.
- Chethan, S., & Malleshi, N. G. (2007). Finger millet polyphenols: Characterization and their nutraceutical potential. *American Journal of Food Technology*, 2(7), 582-592.
- CSA (Central Statistical Agency), (2019). Agricultural sample survey 2018/2019(2011 E.C), report on area and production of major crops private peasant holdings, meher season). Central Statistical Agency (CSA), Volume I, *Statistical Bulletin* 589, June, 2019, Addis Ababa, Ethiopia.
- Damtie, Y., Girma, F., Terfessa, A., & Demelash, H. (2019). Genetic Diversity and Heritability Estimates among Ethiopian Finger Millet (*Eleusine coracana*(L.) Gaertn.) Genotypes for Yield and Its Contributing Traits at Assosa, Western Ethiopia. *Asian Journal of Plant Science and Research*, 9(2), 6-15.
- Falconer, D.S. & F.C. Mackay, (1996). Introduction to Quantitative Genetics. Longman, New York. 464p.
- FAOSTAT (Food and Agriculture Organization of the United Nations Statics division). (2015). Evolution of reproductive behavior of the genus *Eleusine*. *Euphytica*, 31: 399-404.
- Gomez, A.K. & A.A. Gomez, 1984. Statistical Procedure for Agricultural Research. Second Edition. Wiley International Science, New York.
- Johnson, H. W., Robinson, H. F., & Comstock, R. E. (1955). Genotypic and phenotypic correlations in soybeans and their implications in selection 1. *Agronomy journal*, 47(10), 477-483.
- Katiyar, R. P., Prasad, J., Singh, A. B., & Ram, K. (1977). Association analysis of grain yield and its components in segregating populations of chickpea [India]. *Indian Journal of Agricultural Sciences (India)*.
- Keerthana, K., Chitra, S., Subramanian, A., Nithila, S., & Elangovan, M. (2019). Studies on genetic variability in finger millet [*Eleusine coracana* (L.) Gaertn] genotypes under sodic conditions. *Electronic Journal of Plant Breeding*, 10(2), 566-569.
- Kumar, D., Tyagi, V., Ramesh, B., & Pal, S. (2010). Genetic diversity in finger millet (*Eleusine coracana* L.). *Crop improvement*, 37(1), 25-28.
- Patel, S. N., Patil, H. E., & Papat, R. C. (2017). Genetic diversity study in finger millet (*Eleusine coracana* L.) genotypes: A multivariate analysis approach. *Int. J. Pure App. Biosci*, 5(5), 183-189.
- Rudin, B., H.E. Shashidhar, R.S. Kulkarni, S.Hittalmani, (2004). Genetic diversity assessment of finger millet, (*Eleusine coracana* (L.) Gaertn), germplasm through RAPD analysis. *PGR Newsletter* 138: 50-54.
- Patil, H. E., Patel, B. K., & Patel, S. N. (2017). Assessment of genetic diversity in Finger millet (*Eleusine coracana* L.) through multivariate analysis approach. *International Journal of Economic Plants*, 4(4), 148-151.
- IBPGR, (1985). Descriptors for Finger millet. IBPGR secretariat, FAO, Rome, Italy: 1-20.
- Singamsetti, A., Patro, T. S. S. K., Anuradha, N., & Divya, M. (2018). Studies on Genetic Variability for Yield and Yield Attributing Traits in Finger millet (*Eleusine coracana* L. Gaertn). *International Journal of Current Microbiology and Applied Sciences*, 7, 90-95.
- Singh, R. K. (1985). Biometrical methods in Quantitative Genetic Analysis. Kalyani Pub. Ludhiana. *New Delhi, Revised Ed.*, 318.
- Tesfaye, K., & Mengistu, S. (2017). Phenotypic characterization of Ethiopian finger millet accessions (*Eleusine coracana* (L.) Gaertn), for their agronomically important traits. *Acta Universitatis Sapientiae, Agriculture and Environment*, 9(1), 107-118.